

Scaling limits of lattice quantum fields by wavelets

Vincenzo Morinelli^{1,*}, Gerardo Morsella^{1,△}, Alexander Stottmeister^{2,◦}, Yoh Tanimoto^{1,•}

¹ Dipartimento di Matematica, Università di Roma Tor Vergata
Via della Ricerca Scientifica 1, I-00133 Roma, Italy

² Institut für Theoretische Physik, Leibniz Universität Hannover
Appelstraße 2, 30167 Hannover, Germany

October 22, 2020

Abstract

We present a rigorous renormalization group scheme for lattice quantum field theories in terms of operator algebras. The renormalization group is considered as an inductive system of scaling maps between lattice field algebras. We construct scaling maps for scalar lattice fields using Daubechies' wavelets, and show that the inductive limit of free lattice ground states exists and the limit state extends to the familiar massive continuum free field, with the continuum action of spacetime translations. In particular, lattice fields are identified with the continuum field smeared with Daubechies' scaling functions. We compare our scaling maps with other renormalization schemes and their features, such as the momentum shell method or block-spin transformations.

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 2 |
| 2 | Preliminaries | 6 |
| 2.1 | An operator-algebraic renormalization group scheme | 6 |
| 2.2 | Inductive limit of C^* -algebras and representations | 9 |
| 2.3 | Lattice models and local algebras | 10 |
| 2.4 | Continuum field theory | 11 |
| 3 | Wavelet scaling of lattice scalar fields | 12 |
| 3.1 | Scalar fields on lattice | 12 |
| 3.1.1 | First and second quantization | 12 |
| 3.1.2 | Scalar fields on different scales | 13 |
| 3.1.3 | Scaling maps from one-particle spaces | 14 |
| 3.2 | An instructive example: The block-spin method | 15 |
| 3.3 | A generalization: The wavelet method | 16 |
| 3.3.1 | Scaling map from scaling function. | 16 |
| 3.3.2 | Wavelet bases. | 19 |
| 3.3.3 | Block-spin renormalization in terms of orthogonal Haar wavelets | 20 |
| 3.3.4 | The Daubechies wavelets | 21 |

* morinell@mat.uniroma2.it; △ morsella@mat.uniroma2.it;
◦ alexander.stottmeister@itp.uni-hannover.de; • hoyt@mat.uniroma2.it

| | | |
|----------|--|-----------|
| 4 | The continuum limit of the free vacua | 21 |
| 4.1 | States on lattice and continuum fields | 22 |
| 4.1.1 | Ground states of lattice free fields | 22 |
| 4.1.2 | Continuum free scalar field | 23 |
| 4.2 | Constructing the continuum limit | 25 |
| 4.2.1 | Embedding the lattice algebras into the continuum algebra | 26 |
| 4.2.2 | Scaling limits of free-field ground states | 27 |
| 4.2.3 | Density of the wavelet observables in the continuum limit | 31 |
| 4.2.4 | Limit of dynamics | 33 |
| 4.2.5 | Infinite volume limit | 39 |
| 5 | Other scaling-map schemes | 45 |
| 5.1 | The block-spin transformation case | 45 |
| 5.2 | Real-space renormalization: Point-like localization | 47 |
| 5.3 | Momentum-space renormalization: Sharp momentum cutoff | 49 |
| 6 | Consequences and Outlook | 53 |
| 6.1 | Other continuum models and locality from lattices | 54 |
| 6.2 | Extension to fermions | 55 |
| 6.3 | Connections with multi-scale entanglement renormalization (MERA) | 56 |
| 6.4 | Scaling maps and biorthogonal wavelets | 57 |
| 6.5 | Entropy and Type I approximation of local algebras | 57 |
| 6.6 | UV vs. IR scaling limits: Momentum transfer to small scales | 58 |

1 Introduction

The Wilson-Kadanoff renormalization group [47, 80, 81] is a cornerstone in the understanding of classical and quantum many-body system. It provides a conceptual framework that unifies the theory of critical phenomena and universality in (quantum) statistical mechanics with quantum field theory via the existence of infrared fixed points under scale-changing operations. Moreover, the construction of continuum models from lattice approximations via scaling limits has been developed to a wide extend in a mathematically rigorous form in constructive quantum field theory, especially in the classical probabilist framework of the Euclidean approach [39, 45, 32]. While in these works interacting models have been rigorously constructed, the field operators and observables of the lattice and continuum theories are related only indirectly in terms of correlation functions and an analytic continuation of the latter together with a Streater-Wightman reconstruction is necessary to access the operator-algebraic content of the quantum field theory. In the present paper, we start from a lattice approximation to quantum field theories closer in spirit to the earlier constructive works by Friedrichs, Glimm-Jaffe and others [39, 35] and the setting of quantum statistical mechanics. We explicitly implement an operator-algebraic approach [14, 13] to the Wilson-Kadanoff renormalization group for scalar lattice quantum fields in any dimension (see also [73] for a short overview). Specifically, using results of wavelet theory [23, 58], we build a real-space renormalization group by identifying the lattice fields with the continuum fields smeared with suitable functions which generate a wavelet basis. This results in a sequence of homomorphisms between lattice field algebras at various scales. We supplement the lattice algebras with a sequence of states that approaches a critical point in an appropriate way (the massless free field for the massive free field), which assures the continuity of asymptotic

symmetry groups avoiding certain obstacles encountered in [46, 50, 66], while keeping track of local algebras, cf. [86]. Therefore, we succeed to construct a continuum field theory in the Haag-Kastler sense [41] in the real-space framework of the Wilson-Kadanoff renormalization group.

Our implementation is inspired by the strong connection between renormalization of classical lattice systems and wavelet theory [4], which prevails in the quantum setting, cp. [12, 28, 57, 36, 84, 83]. As an obvious application, we show rigorously and in detail that the renormalization group flow of the ground states of free lattice fields in the vicinity of the unstable, massless fixed point allows us to reconstruct the massive continuum free field as a scaling limit. By using compactly supported wavelets (or finite low-pass filters), this construction directly yields local time-zero algebras of the continuum theory which allows us to pass to the infinite-volume limit in a simple way. From a mathematical point of view, operator algebras naturally accommodate the quantum version of the Wilson-Kadanoff approach as the former are generally understood to incorporate non-commutative measure theory, see [51, 74] for another approach reversing the classical Euclidean approach to obtain a quantum version. Furthermore, in the specific formulation we use, the renormalization group is realized as an inductive system of $*$ -morphisms in our examples, and the scaling limit is identified with a state on the inductive-limit algebra [48]. To some extent, this reflects a weak form of universality [87, Chapter 3] in mathematical sense: The scaling limit is independent of the specific details of the inductive system at finite scales, e.g. modifications of the dispersion relations of the free lattice ground states are irrelevant as long as the asymptotic behavior is unchanged. As both, the infrared limits of the renormalization group and the inductive limits of operators algebras, are only concerned with asymptotic properties of their building blocks (with a certain notion of coherence), this suggests a link at a conceptual level. This should not be confused with the concept of universality classes which are associated with different fixed points of the renormalization group when taking into account additional interactions. In the latter sense, we are only concerned with the universality class of the massless free field in this work. As can be seen from the free field example, criticality of the lattice models (i.e. divergence of correlation lengths) is intimately linked to continuity properties of the operator-algebraic limits.

Our formulation of the renormalization group is also linked to other developments in quantum theoretical descriptions of low-dimensional many-body systems, e.g. the density matrix renormalization group (DMRG) [79, 77, 78, 70], which is a real-space formulation that has turned out to be especially efficient in one dimension. As the Wilson-Kadanoff approach fits naturally into an information geometrical setup [5], real-space renormalization schemes have also drawn increasing attention in the theory of quantum information, specifically concerning tensor networks [18] and the multi-scale entanglement renormalization ansatz (MERA) [75, 27, 28, 67].

Summary of the formulation and results

We formulate a renormalization group for scalar lattice field theories using a scaling function ϕ of wavelet theory. Such a function satisfies a self-similarity equation,

$$\phi(x) = \sum_{n \in \mathbb{Z}^d} h_n 2^{\frac{d}{2}} \phi(2x - n),$$

which encodes the decomposition of ϕ at coarse scale in terms of itself at a fine scale. Clearly, this equation is analogous in spirit to Kadanoff's block-spin transformation, which considers a spin on a coarse lattice to be an average of spins on a fine lattice. But, while the block-spin

transformation typically averages uniformly over adjacent sites, the scaling equation allows for a weighted average over sites. The coefficients h_n are commonly known as a low-pass filter as they describe the coarse-scale features captured by ϕ . Wavelets enter the picture as a parametrization of the L^2 -complement of ϕ after increasing the resolution leading to a complementary high-pass filter tracking the fine-scale details. Thus, it is tempting to utilize the scaling equation to define the relation among lattice quantum fields at successive scales ε_N , $\varepsilon_{N+1} = \varepsilon_N/2$ (and similarly for momenta):

$$\Phi_N(x) = 2^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n \Phi_{N+1}(x + \varepsilon_{N+1}n).$$

In terms of the algebras of observables \mathfrak{A}_N at scales ε_N this precisely defines an inductive system,

$$\alpha_{N'}^N : \mathfrak{A}_N \longrightarrow \mathfrak{A}_{N'}, \quad N < N',$$

with the algebraic and analytic properties of the α 's encoded into the coefficients h_n . Specifically, by using a Daubechies' scaling function which is compactly supported and orthogonal (integer translates of ϕ are orthogonal at a fixed scale), it is possible to realize the α 's as unital $*$ -morphisms that equip the limit algebra $\mathfrak{A}_\infty = \varinjlim_N \mathfrak{A}_N$ with a natural quasi-local structure because the low-pass filter is finite in this case and, therefore, limits the spatial support of a localized finite-scale quantum field $\Phi_N(x)$ in the limit. Indeed, the image of the latter in \mathfrak{A}_∞ can be identified with the continuum quantum field Φ smeared with ϕ at scale ε_N ,

$$\Phi_N(x) = \varepsilon_N^{-d} \int dy \Phi(y) \phi(\varepsilon_N^{-1}(y - x)).$$

Now, the α 's are what we will call the renormalization group or scaling maps, and the more common picture of the renormalization group transformations linking states of a physical system among different scales arises by their dual action on the states spaces $\mathfrak{S}_{N+1} \rightarrow \mathfrak{S}_N$ (generalized density matrices). Following Wilson [82], we consider the renormalization group flow of initial states $\omega_0^{(N)}$ of the finite scale systems \mathfrak{A}_N ,

$$\omega_M^{(N)} = \omega_0^{(N+M)} \circ \alpha_{N+M}^N,$$

to construct a scaling limit $\omega_\infty^{(\infty)} = \varprojlim_N \lim_M \omega_M^{(N)}$ on \mathfrak{A}_∞ . To make these abstract considerations maximally explicit and to test the construction, we provide a rather complete treatment of the important case of harmonic (free) lattice fields on the torus L^d ,

$$H_0^{(N)} = \frac{1}{2} \varepsilon_N^d \left(\sum_{x \in \Lambda_N} \Pi_{N|x}^2 + \mu_N^2 \varepsilon_N^{-2} \Phi_{N|x}^2 - 2 \sum_{\langle x,y \rangle \subset \Lambda_N} \varepsilon_N^{-2} \Phi_{N|x} \Phi_{N|y} \right),$$

and their ground states $\omega_L^{(N)}$, which are expected to admit sensible scaling limits in the vicinity of the Gaussian fixed point $\mu_N = 2d$. More precisely, we show that:

- (1) the scaling limit $\omega_{L,\infty}^{(\infty)}$ exists in finite volume $(2L)^d$ and is given by the continuum free field of mass m assuming the renormalization condition $\lim_{N \rightarrow \infty} \varepsilon_N^{-2} (\mu_N^2 - 2d) = m^2$ holds,

- (2) the finite scale dynamics converges to the free dynamics in the Gelfand-Naimark-Segal representation of $\omega_{L,\infty}^{(\infty)}$ (a similar statement holds for spatial translations),
- (3) the infinite volume (or thermodynamical) limit $L \rightarrow \infty$ of $\omega_{L,\infty}^{(\infty)}$ exists and the local time-zero and spacetime algebras in finite volume are unitarily equivalent to those in infinite volume,

if the renormalization group is realized in terms of Daubechies' wavelets of sufficiently high regularity. The latter observation also fits nicely with the idea of universality as the details of a valid approximation should not matter, i.e. the scaling limit is independent of the precise regularity of wavelets as long as it is sufficient. Thus, our operator-algebraic implementation of Wilson-Kadanoff renormalization enables a direct construction of the local net of observables of the continuum free field (in the sense of the Haag-Kastler axioms [41]) from free lattice fields. In addition, we observe that our approach yields a rigorous way to deduce spacetime locality in the continuum from Lieb-Robinson bounds [52, 21, 65, 62, 64] albeit asymptotically optimal estimates of the Lieb-Robinson velocity are required to conclude that $c = 1$ in natural units. Moreover, we also illustrate how our wavelet renormalization group yields an analytic versions of MERA for scalar fields in arbitrary dimensions, cp. [30, 31, 42, 83]. Finally, we also comment on the approach to the renormalization group via scaling algebras [15, 16, 22, 9], where the principal difference to our work lies in the consideration of the ultraviolet limit of theories already defined in the continuum rather than of the infrared limit of lattice theories.

The article is structured as follows. In Section 2, we recall the operator-algebraic formulation of Wilson-Kadanoff renormalization, and we introduce the required objects for the construction of scaling limits such as scaling maps, limit algebras and scaling limits of states and dynamics. We also introduce a notion of spatial local algebras for lattice models and give a tentative definition of a continuum theory in terms of local observables in finite and infinite volume that reasonable scaling limit should fit into. In Section 3 we invoke the framework of second quantization for the description of scalar fields at a fixed scale, and wavelet scaling maps are introduced as a generalization of the block-spin transformation, which we complement by essential ingredients from the theory of wavelets. In Section 4, we construct the scaling limit of free, massive lattice scalar fields and show that the limit state gives the massive free state while the lattice fields are represented as the continuum field smeared with the scaling function. Section 5, we discuss the block-spin transformation from the perspective of the wavelet scaling map, and we explain how the former leads to a continuum limit albeit in a more singular sense. Furthermore, we illustrate two other commonly used renormalization groups, point-like localizations and sharp momentum cut-offs (momentum shells), and we show that the first leads to singular limits similar to the block-spin case, while the second is well-adapted to dynamics but intrinsically non-local. Section 6 outlines various consequences of the previous sections thereby giving an outlook on topics for future research.

Acknowledgements

AS would like to thank Tobias J. Osborne and Reinhard F. Werner for useful discussions and comments. VM thanks Domenico Marinucci and Claudio Durastanti for comments and discussions. AS was in part supported by Alexander-von-Humboldt Foundation through a Feodor Lynen Return Fellowship. The European Research Council Advanced Grant 669240 QUEST supported VM and partially GM. VM was supported by Indam from March 2019 to February 2020. Until February 2020 YT was supported by Programma per giovani

ricercatori, anno 2014 “Rita Levi Montalcini” of the Italian Ministry of Education, University and Research. VM, GM and YT also acknowledge the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome “Tor Vergata”, CUP E83C18000100006 and the University of Rome “Tor Vergata” funding scheme “Beyond Borders”, CUP E84I19002200005.

2 Preliminaries

2.1 An operator-algebraic renormalization group scheme

The Wilson-Kadanoff renormalization group. A widely used scheme to analyze continuum limits using effective lattice models is the Wilson-Kadanoff renormalization group [47, 82, 76]. For a family of lattices $\{\Lambda_N\}_{N \in \mathbb{N}_0}$, say in \mathbb{R}^d or in the torus \mathbb{T}_L^d of side length $2L$, we consider a sequence of Hamiltonian quantum systems $\{\mathfrak{A}_N, \mathcal{H}_N, H_0^{(N)}\}_{N \in \mathbb{N}_0}$ indexed by the level $N \in \mathbb{N}_0$, the logarithmic scale accounting for the relative density of lattice points (increasing with N). At each level N , $\mathfrak{A}_N \subset \mathcal{B}(\mathcal{H}_N)$ is a concrete C^* -algebra of observables acting on a Hilbert space \mathcal{H}_N , and $H_0^{(N)}$ is an initial Hamiltonian with domain $\mathcal{D}_N \subset \mathcal{B}(\mathcal{H}_N)$. The essential equation of renormalization group theory defines the renormalized Hamiltonians $H_M^{(N)}$ for $N' - N = M \geq 0$ by demanding the equality of partition functions of $H_M^{(N)}$ and $H_0^{(N+M)}$ ([33], see also [34, 77, 78]):

$$Z_0^{(N+M)} = \text{Tr}_{N+M} \left(e^{-H_0^{(N+M)}} \right) = \text{Tr}_N \left(e^{-H_M^{(N)}} \right) = Z_M^{(N)}, \quad (2.1)$$

where both exponentials are assumed to be trace class. More precisely, the renormalization group should define coarse-graining transformations (or quantum operations) between the spaces of density matrices, \mathcal{S}_{N+M}^1 and \mathcal{S}_N^1 , on \mathfrak{A}_{N+M} respectively \mathfrak{A}_N , such that:

$$\mathcal{E}_N^{N+M} \left(\rho_0^{(N+M)} \right) = \rho_M^{(N)}, \quad (2.2)$$

with $\rho_M^{(N)} = (Z_M^{(N)})^{-1} e^{-H_M^{(N)}}$. The transformation are also know as the descending superoperators [27]. Since the choice of levels $N < N' = N + M$ is arbitrary, we naturally require the semi-group property:

$$\mathcal{E}_N^{N'} \circ \mathcal{E}_{N'}^{N''} = \mathcal{E}_N^{N''}, \quad N < N' < N''. \quad (2.3)$$

At the level of Hamiltonians with corresponding maps, $H_0^{(N+M)} \mapsto H_M^{(N)}$, this is summarized by Wilson’s *triangle of renormalization* [82, p. 790] in Figure 1.

Operator-algebraic renormalization. Let us translate (2.2) into a statement involving the algebras $\{\mathfrak{A}_N\}_{N \in \mathbb{N}_0}$ and states on them (generalizing density matrices). We find for $a_N \in \mathfrak{A}_N$:

$$\text{Tr}_N \left(\rho_M^{(N)} a_N \right) = \text{Tr}_N \left(\mathcal{E}_N^{N+M} \left(\rho_0^{(N+M)} \right) a_N \right) = \text{Tr}_{N+M} \left(\rho_0^{(N+M)} \alpha_{N+M}^N(a_N) \right) \quad (2.4)$$

Here, $\alpha_{N+M}^N : \mathfrak{A}_N \rightarrow \mathfrak{A}_{N+M}$ is the dual of \mathcal{E}_N^{N+M} (or an ascending superoperators as in [27]). Thus, for a given family of initial states $\{\omega_0^{(N)}\}_{N \in \mathbb{N}_0}$ instead of Hamiltonian $\{H_0^{(N)}\}_{N \in \mathbb{N}_0}$, (2.2) generalizes to:

$$\omega_0^{(N+M)} \circ \alpha_{N+M}^N = \omega_M^{(N)}. \quad (2.5)$$

$$\begin{array}{ccccccc}
& & & & & & H_0^{(N+2)} \longrightarrow \dots \\
& & & & & & \downarrow \\
& & & & & & H_0^{(N+1)} \longrightarrow H_1^{(N+1)} \longrightarrow \dots \\
& & & & & & \downarrow \quad \downarrow \\
& & & & & & H_0^{(N)} \longrightarrow H_1^{(N)} \longrightarrow H_2^{(N)} \longrightarrow \dots
\end{array}$$

Figure 1: Wilson's triangle of renormalization in terms of Hamiltonians of subsequent scales. The horizontal arrows arise from a choice of the initial Hamiltonians $H_0^{(N)}$ and their coarse-graining $H_0^{(N+M)} \rightarrow H_M^{(N)}$.

A natural requirement is that α_{N+M}^N is unital and completely positive because it should map states into states and preserve probability as expressed by the equality of partition functions (2.1). We further require that $\alpha_{N'}^N$ is a $*$ -morphism, so that we can define the inductive limit algebra. The semi-group property (2.3) translates to:

$$\alpha_{N''}^{N'} \circ \alpha_{N'}^N = \alpha_{N''}^N. \quad (2.6)$$

We call the collection $\{\alpha_{N'}^N\}_{N < N' \in \mathbb{N}_0}$, the *scaling maps* or *renormalization group*. At the level of algebras and states, Figure 2 provides an analogue to Figure 1:

$$\begin{array}{ccccccc}
& & & \dots & & \vdots & \vdots \\
& & & \omega_0^{(N+2)} \longrightarrow \dots & & \omega_\infty^{(N+2)} & \mathfrak{A}_{N+2} \\
& & & \mathcal{E}_{N+1}^{N+2} \downarrow & & & \uparrow \alpha_{N+2}^{N+1} \\
& & & \omega_0^{(N+1)} \longrightarrow \omega_1^{(N+1)} \longrightarrow \dots & & \omega_\infty^{(N+1)} & \mathfrak{A}_{N+1} \\
& & & \mathcal{E}_N^{N+1} \downarrow \quad \mathcal{E}_N^{N+1} \downarrow & & & \uparrow \alpha_{N+1}^N \\
& & & \omega_0^{(N)} \longrightarrow \omega_1^{(N)} \longrightarrow \omega_2^{(N)} \longrightarrow \dots & & \omega_\infty^{(N)} & \mathfrak{A}_N
\end{array}$$

Figure 2: Wilson's triangle of renormalization in terms of algebras and states at subsequent scales.

Example 2.1. A simple example of quantum operations $\{\mathcal{E}_N^{N'}\}_{N < N' \in \mathbb{N}_0}$ are partial traces $\text{Tr}_{N' \rightarrow N}$ for $\mathcal{H}_N = \mathcal{H}_0^{\otimes 2^N}$ and $\mathfrak{A}_N = B(\mathcal{H}_N)$ such that $\alpha_{N'}^N(a_N) = a_N \otimes \mathbf{1}_{N'-N}$. Natural generalizations of the latter arising in the context of lattice gauge theory are [59, 14, 13]:

$$\alpha_{N'}^N(a_N) = U_{N'}(a_N \otimes \mathbf{1}_{N'-N})U_{N'}^*, \quad (2.7)$$

for some unitary $U_{N'} \in \mathcal{U}(\mathcal{H}_{N'})$.

Equation (2.7) is satisfied for our construction of a renormalization group for lattice scalar fields and is analogous to MERA (multi-scale entanglement renormalization), see (6.5).

Continuum limit, inductive limit algebra and scaling limit state. Our operator-algebraic renormalization provides an algorithm for the construction of a continuum limit of lattice models, which we roughly split into three parts: First, we construct a family of scaling maps $\{\alpha_{N+1}^N\}_{N \in \mathbb{N}_0}$ between algebras of lattice fields to define the renormalization group arriving at an inductive limit algebra (see Section 2.2 for a summary of the construction):

$$\mathfrak{A}_\infty = \varinjlim_{N \in \mathbb{N}_0} \mathfrak{A}_N. \quad (2.8)$$

Second, we consider a sequence of initial states $\{\omega_0^{(N)}\}_{N \in \mathbb{N}_0}$ (at the upside of the triangle in Figure 2). Their restrictions to coarser lattices are determined by the renormalization group elements $\alpha_{N'}^N$ (generalizing the coarse-graining transformations $\{\mathcal{E}_{N'}^N\}$ via pullback) according to (2.5). On each lattice algebra \mathfrak{A}_N , the sequence $\{\omega_M^{(N)}\}_{M \in \mathbb{N}_0}$ (horizontal lines) should admit a limit state,

$$\omega_\infty^{(N)} = \lim_{M \rightarrow \infty} \omega_M^{(N)}, \quad (2.9)$$

at least for a subsequence, and we expect the limit state to be stable under coarse graining,

$$\omega_\infty^{(N')} \circ \alpha_{N'}^N = \omega_\infty^{(N)}, \quad (2.10)$$

because formally $\alpha_\infty^{N'} \circ \alpha_{N'}^N = \alpha_\infty^N$ for $\alpha_\infty^N = \lim_{N' \rightarrow \infty} \alpha_{N'}^N$. Algebraically, the consistency expressed by this stability condition would allow for the existence of a projective-limit state on the inductive-limit algebra \mathfrak{A}_∞ (cf. Section 2.2),

$$\omega_\infty^{(\infty)} = \varprojlim_{N \in \mathbb{N}_0} \omega_\infty^{(N)}. \quad (2.11)$$

Such projective-limit states will be called *scaling limits* of the initial sequence $\{\omega_0^{(N)}\}_{N \in \mathbb{N}_0}$, and their existence together with the property (2.10) is assured under rather mild conditions as shown in Proposition 2.2 below. The expected non-uniqueness of (2.9) and, therefore, of the scaling limits is physically meaningful (e.g. renormalization group trajectories and phase transitions) and is partially reflected in the need for *renormalization conditions*, e.g. on the couplings implicitly present in the initial states.

Finally given a scaling limit $\omega_\infty^{(\infty)}$ together with the algebra \mathfrak{A}_∞ , we perform the Gelfand-Naimark-Segal (GNS) construction to arrive at a Hilbert space representation,

$$\pi_\infty^{(\infty)} : \mathfrak{A}_\infty \longrightarrow \mathcal{H}_\infty^{(\infty)}, \quad \omega_\infty^{(\infty)}(a) = \langle \Omega_\infty, \pi_\infty^{(\infty)}(a) \Omega_\infty \rangle_{\mathcal{H}_\infty^{(\infty)}}, \quad (2.12)$$

such that the scaling limit is implemented by the vector state $\Omega_\infty \in \mathcal{H}_\infty^{(\infty)}$. Now, $\mathcal{H}_\infty^{(\infty)}$ should be regarded as the Hilbert space of the continuum limit, we can identify the elements of each \mathfrak{A}_N with certain operators of the continuum field via scaling maps as expressed by (2.8)) and compare the expected properties. As we are working in a Hamiltonian setting, the continuum limit should be interpreted in terms of time-zero fields. Moreover, natural candidates for non-trivial scaling limits are families of ground states $\{\omega_{\lambda_N}\}_{N \in \mathbb{N}_0}$ for lattice Hamiltonians $\{H_{\lambda_N}\}_{N \in \mathbb{N}_0}$ admitting quantum critical points¹ (not necessarily quantum phase transitions) with respect to their respective (dimensionless) couplings $\{\lambda_N\}_{N \in \mathbb{N}_0}$, see e.g. [69]. In the case of harmonic lattice fields, we will see that families of lattice ground states admit the vacuum states of arbitrary masses (reflecting the non-uniqueness) of the continuum free field as scaling limits.

¹Physically, a necessary condition for the existence of a non-trivial sequence of limit states $\{\omega_\infty^{(N)}\}_{N \in \mathbb{N}_0}$ is the divergence (in units of lattice length) of the sequence of correlation lengths $\{\xi_{N'}^{(N)}\}_{N' \in \mathbb{N}_0}$ associated with pairs of sequences $\{\alpha_{N'}^N(a_N), \alpha_{N'}^N(b_N)\}_{N' \in \mathbb{N}_0}$ of local operators for any $N' \in \mathbb{N}_0$.

Dynamics. Besides the existence of a scaling limit $\{\omega_\infty^{(\infty)}\}$, we can also analyze the convergence of the family of time-evolution groups $\{\eta_t^{(N)} = e^{itH_0^{(N)}}(\cdot)e^{-itH_0^{(N)}}\}_{N \in \mathbb{N}_0}$. In view of the similarities of our renormalization group scheme with the construction of thermodynamical limits in quantum statistical mechanics, we consider, for this purpose, sequences of the form,

$$\{\alpha_\infty^{N'}(\eta_t^{(N')}(a_N))\}_{N' > N}, \quad (2.13)$$

for fixed $t \in \mathbb{R}$ and $a_N \in \mathfrak{A}_N$ and ask whether these are Cauchy sequences in the Hilbert space representation relative to the scaling limit $\omega_\infty^{(\infty)}$, i.e. for a suitable operator topology on $\pi_\infty^{(\infty)}(\mathfrak{A}_\infty)$ relative to the scaling limit [41, 11, 63]. This way we may define the limit $\eta_t^{(\infty)} = \lim_{N \rightarrow \infty} \eta_t^{(N)}$ on the closure $\mathcal{A} = \overline{\pi_\infty^{(\infty)}(\mathfrak{A}_\infty)}$ and obtain a scaling-limit Hamiltonian $H_\infty^{(\infty)}$ (corresponding to an extrapolation of the upside of Wilson's triangle in Figure 1) given suitable continuity properties of $\eta^{(\infty)} : \mathbb{R} \curvearrowright \mathcal{A}$.

Free field as an example. For general interacting lattice models, we do not expect to find all the above objects in closed form, but suitable expansion or perturbation methods will be required to obtain approximations [8]. Moreover, the extent to which it will be possible to carry out this formulation of Wilson-Kadanoff renormalization will depend sensitively on the choice of scaling maps $\alpha_{N'}^N$, and the amount of control over the state space $\mathfrak{S}(\mathfrak{A}_N) = \mathfrak{S}_N$ and the effective Hamiltonian $H_0^{(N)}$ of each lattice system $(\mathfrak{A}_N, \mathcal{H}_N)$. Nevertheless, we show in the following that this scheme can be carried out in real space for the free scalar field in any dimension with full control over all involved objects.

2.2 Inductive limit of C^* -algebras and representations

The construction of a sequence of lattice C^* -algebras and its inductive limit is central to our construction of a continuum limit. Therefore, we recall this fundamental architecture to present detailed and self-contained understanding of the limit procedure. For further references see for instance [48, 56, 7]

Following the notation of the previous section, let $\{\mathfrak{A}_N\}_{N \in \mathbb{N}_0}$ be a sequence of unital C^* -algebras. For $N < N' < N''$, we assume that there exists a unital injective $*$ -morphism $\alpha_{N'}^N$ from \mathfrak{A}_N to $\mathfrak{A}_{N'}$ such that $\alpha_{N''}^{N'} \circ \alpha_{N'}^N = \alpha_{N''}^N$. Since $\alpha_{N'}^N$ is injective, it is also isometric. $\{\mathfrak{A}_N\}_{N \in \mathbb{N}_0}$ together with $\{\alpha_{N'}^N : N, N' \in \mathbb{N}_0, N < N'\}$ is called a *directed system of C^* -algebras*. Such a directed system of C^* -algebras can be embedded up to $*$ -isomorphisms into a C^* -algebra called the *inductive limit*. In particular, there exists a C^* -algebra \mathfrak{A}_∞ such that:

- (i) for every $N \in \mathbb{N}_0$ there exists a unital injective $*$ -homomorphism α_∞^N from \mathfrak{A}_N to \mathfrak{A}_∞ . Furthermore, if $N < N'$ then $\alpha_\infty^N = \alpha_\infty^{N'} \circ \alpha_{N'}^N$ and $\bigcup_{N \in \mathbb{N}_0} \alpha_\infty^N(\mathfrak{A}_N)$ is dense in \mathfrak{A}_∞ ;
- (ii) the C^* -algebra \mathfrak{A}_∞ in (i) is unique up to a $*$ -isomorphism; in particular, if \mathfrak{B}_∞ is another C^* -algebra with $*$ -homomorphisms β_∞^N satisfying conditions as in (i), then there exists a $*$ -isomorphism $\mathcal{J} : \mathfrak{B}_\infty \rightarrow \mathfrak{A}_\infty$ such that $\alpha_\infty^N = \mathcal{J} \circ \beta_\infty^N$.

Next, we consider a family of states $\{\omega^{(N)}\}_{N \in \mathbb{N}_0}$ such that $\omega^{(N)}$ is a state on \mathfrak{A}_N , and we assume that $\omega^{(N)} = \omega^{(N')} \circ \alpha_{N'}^N$. Such a family is said to be *projectively consistent*. Then, there is a uniquely defined state $\omega^{(\infty)}$ on \mathfrak{A}_∞ such that $\omega^{(\infty)} \circ \alpha_\infty^N = \omega^{(N)}$. $\omega^{(\infty)} = \varprojlim_{N \in \mathbb{N}_0} \omega^{(N)}$ is called the *projective limit* of the family $\{\omega^{(N)}\}$.

As our notion of continuum limit from the previous section is intimately linked with the GNS construction, we also recall the following: The GNS representation $(\pi^{(\infty)}, \mathcal{H}^{(\infty)}, \Omega_\infty)$ of $(\mathfrak{A}_\infty, \omega^{(\infty)})$ extends the GNS representation $(\pi^{(N)}, \mathcal{H}^{(N)}, \Omega_N)$ associated with $(\mathfrak{A}_N, \omega^{(N)})$. More precisely, the GNS construction induces unique isometries $V_{N'}^N$ from $\mathcal{H}^{(N)}$ into $\mathcal{H}^{(N')}$ such that $\pi^{(N')}(\alpha_{N'}^N(a_N))V_{N'}^N = V_{N'}^N\pi^{(N)}(a_N)$ for all $a_N \in \mathfrak{A}_N$ and $V_{N'}^N\Omega_N = \Omega_{N'}$. Therefore, the GNS representations $(\pi^{(N)}, \mathcal{H}^{(N)}, \Omega_N)$ naturally acquire the structure of a compatible direct system of Hilbert spaces, which admits an inductive limit given by $(\pi^{(\infty)}, \mathcal{H}^{(\infty)}, \Omega_\infty)$ together with isometries V_∞^N from $\mathcal{H}^{(N)}$ into $\mathcal{H}^{(\infty)}$ such that $\pi^{(\infty)}(\alpha_\infty^N(a_N))V_\infty^N = V_\infty^N\pi^{(N)}(a_N)$ for all $a_N \in \mathfrak{A}_N$ and $V_\infty^N\Omega_N = \Omega_\infty$. The unique normal extension of $\omega^{(\infty)}$ to the von Neumann algebra $\pi^{(\infty)}(\mathfrak{A}_\infty)''$ will be denoted by the same symbol.

In view of the construction and existence of projectively consistent families of states (cf. (2.9) and (2.11)), we make the important observation (which generalizes to directed index sets):

Proposition 2.2. *Let $\{\mathfrak{A}_N\}_{N \in \mathbb{N}_0}$, $\{\alpha_{N'}^N\}_{N < N'}$ be a directed system of C^* -algebras, and let $\{\omega^{(N)}\}_{N \in \mathbb{N}_0}$ be a family of states on $\{\mathfrak{A}_N\}_{N \in \mathbb{N}_0}$. For $M > 0$, we define a state by $\omega_M^{(N)} := \omega^{(N+M)} \circ \alpha_{N+M}^N$ on \mathfrak{A}_N . If, for each N , $\lim_{M \rightarrow \infty} \omega_M^{(N)} =: \omega_\infty^{(N)}$ exists in the weak* topology, then it defines a projectively consistent family $\{\omega_\infty^{(N)}\}_{N \in \mathbb{N}_0}$ and hence $\varprojlim_{N \in \mathbb{N}_0} \omega_\infty^{(N)} = \omega_\infty^{(\infty)}$ is well-defined.*

Proof. Because of weak* convergence, we have:

$$\lim_{M \rightarrow \infty} \left| \omega_\infty^{(N')} \circ \alpha_{N'}^N(a_N) - \omega_M^{(N')} \circ \alpha_{N'}^N(a_N) \right| = 0 \quad \text{for all } a_N \in \mathfrak{A}_N$$

for all $N < N'$. Therefore, the following is true for all $N < N'$ (assuming $M > N' - N$):

$$\begin{aligned} \omega_\infty^{(N)} &= \lim_{M \rightarrow \infty} \omega_M^{(N)} = \lim_{M \rightarrow \infty} \omega^{(N+M)} \circ \alpha_{N+M}^N = \lim_{M \rightarrow \infty} \omega^{(N+M)} \circ \alpha_{N+M}^{N'} \circ \alpha_{N'}^N \\ &= \lim_{M \rightarrow \infty} \omega_{N+M-N'}^{(N')} \circ \alpha_{N'}^N = \omega_\infty^{(N')} \circ \alpha_{N'}^N. \end{aligned}$$

□

It may happen that each $\{\omega_M^{(N)}\}_{M \in \mathbb{N}_0}$ converges only by taking a subsequence, and, thus, the limit may not be unique. In the context of scaling limits, this corresponds to the non-uniqueness of the vacuum or different coupling constants and, therefore, is physically important. Nevertheless, the proposition implies the existence of scaling limits in general due to the weak* compactness² of the state space of C^* -algebras.

2.3 Lattice models and local algebras

We are mainly interested in field theories on hypercubic lattices Λ_N and their scaling limits. This means that each \mathfrak{A}_N is equipped with a local structure as follows.

As in Section 2.1, we associate to each N , the scale, a spatial lattice Λ_N with lattice spacing ε_N such that $\varepsilon_{N+1} < \varepsilon_N$. To each lattice point $x \in \Lambda_N$ we assign a local algebra $\mathfrak{A}_N(x)$. The algebras $\mathfrak{A}_N(x)$ and $\mathfrak{A}_N(y)$ should commute if $x \neq y$, and they generate the whole algebra \mathfrak{A}_N . For a subset X of Λ_N , let us denote $\mathfrak{A}_N(X) = \overline{\bigcup_{x \in X} \mathfrak{A}_N(x)}^{\|\cdot\|}$.

Furthermore, we assume that the image of the renormalization map $\alpha_{N+1}^N(\mathfrak{A}_N(x))$ should be included in $\mathfrak{A}_{N+1}(S_x)$, where $S_x = \{y \in \Lambda_{N+1} : y - x \in [0, \varepsilon_{N+1} r_{\max}]^d\}$, and $r_{\max} > 0$

²Apply a diagonal-sequence argument along $N \in \mathbb{N}_0$ to the convergent subsequences $\{\omega_{M_{k_N}}^{(N)}\}_{M_{k_N} > 0}$.




Figure 3: Geometric subdivisions of the lattice Λ_1^L for $d = 2$.

does not depend on N . We call the collection $\{\mathfrak{A}_N, \alpha_{N'}^N, \Lambda_N\}_{N, N' \in \mathbb{N}, N' > N}$ a *lattice field theory*.

In the following, we take standard hypercubic lattices in \mathbb{R}^d with $\varepsilon_N = 2^{-N}\varepsilon$ for some $\varepsilon > 0$, hence $\Lambda_N \subset \Lambda_{N+1}$. However, the renormalization map α_{N+1}^N is in general *not* just the identification of $\alpha_{N+1}^N(\mathfrak{A}_N(x))$ with $\mathfrak{A}_{N+1}(x)$ for $x \in \Lambda_N \subset \Lambda_{N+1}$. Indeed, the key to obtain the continuum field as the scaling limit is to identify a lattice field in \mathfrak{A}_N with a smeared field in the continuum, not a point-like field. This naturally leads to the wavelet scaling in Section 3.3.

By the above, the increase in support due to $\alpha_{N'}^N$ for an element $a_N \in \mathfrak{A}_N$ is bounded from above by $r_{\max}\varepsilon_N \left(1 - 2^{-(N'-N)}\right)$. Thus, we can define local algebras $\mathfrak{A}_\infty(S) \subset \mathfrak{A}_\infty$ for suitable open domains $S \subset \mathbb{T}_L^d = [-L, L]^d$ by collecting at each level N all the operators a_N with support in the sublattice $\Lambda_N(S) \subset \Lambda_N \cap S$ with the convention that the cube $x + [0, \varepsilon_N r_{\max}]^d$ does not intersect the boundary ∂S for each site $x \in \Lambda_N(S)$, see Figure 4. The bound on the increase of support ensures that this definition is compatible with the equivalence classes formed with respect to the inductive system (3.11):

$$\mathfrak{A}_\infty(S) = \varinjlim_{N \in \mathbb{N}_0} \mathfrak{A}_N(\Lambda_N(S)), \quad (2.14)$$

It immediately follows from locality at level N that

$$\mathfrak{A}_\infty(S) \subset \mathfrak{A}_\infty(S') \quad S \subset S', \quad (2.15)$$

$$[\mathfrak{A}_\infty(S), \mathfrak{A}_\infty(S')] = \{0\} \quad S \cap S' = \emptyset. \quad (2.16)$$

We define $\mathfrak{A}_\infty = \overline{\bigcup_S \mathfrak{A}_\infty(S)}$ a quasi-local algebra [10].

2.4 Continuum field theory

Let $\omega_\infty^{(\infty)}$ be the scaling limit state (2.11) of the sequence $\{\omega_0^{(N)}\}$ on a lattice field theory $\{\mathfrak{A}_N, \alpha_{N'}^N, \Lambda_N\}$. In the GNS representation $\pi_\infty^{(\infty)}$ of \mathfrak{A}_∞ with respect to $\omega_\infty^{(\infty)}$, we obtain a family of von Neumann algebras $\pi_\infty^{(\infty)}(\mathfrak{A}_\infty(S))''$. These algebras should be a continuum field theory, in the following sense.

We say that (\mathcal{A}, U, Ω) is a **continuum time-zero net of observables** in \mathcal{H} , where $\mathcal{A}(S)$ is a von Neumann algebra for each region $S \subset \mathbb{T}_L^d$, U is a strongly continuous unitary representation of $\mathbb{T}_L^d \times \mathbb{R}$ and Ω is a vector such that

- (1) If $S_1 \subset S_2 \subset \mathbb{T}_L^d$, then $\mathcal{A}(S_1) \subset \mathcal{A}(S_2)$.
- (2) $\text{Ad } U(a, 0)(\mathcal{A}(S)) = \mathcal{A}(S + a)$ for $a \in \mathbb{T}_L^d$.
- (3) There is $c' > 0$ such that, if $d(S_1, S_2) > \varepsilon c'$ (where $d(S_1, S_2) = \inf_{x \in S_1, y \in S_2, n \in \mathbb{Z}^d} |x - y + 2Ln|$ is the distance on \mathbb{T}_L^d between two regions S_1, S_2), then $\text{Ad } U(0, t)(\mathcal{A}(S_1))$ commutes with $\mathcal{A}(S_2)$ for $t < \varepsilon$.




Figure 4: Illustration of a localized sublattice $\Lambda_N(S)$ (the black sites) for $r_{\max} = 1$ in dimension $d = 2$. Thick lines attached to black sites indicate boundaries of support in the scaling limit.

(4) $U(a, t)\Omega = \Omega$.

(5) The generator of the one-parameter group $U(0, t)$ is positive.

If \mathbb{T}_L^d is replaced by \mathbb{R}^d , we call it a **continuum (infinite volume) time-zero net of observables**. If $c = 1$, these properties are restrictions of the Haag-Kastler axioms [41] to the time-zero plane and the restriction of any Haag-Kastler net to the $t = 0$ plane satisfies them. We do not include uniqueness of the vacuum, because it may fail for a physical reason in the construction through continuum limit (phase transition).

The Poincaré covariance does not follow from these time-zero axioms. Indeed, it is an additional requirement that U extends to a representation of the Poincaré group. A counterexample should be obtained if we start with the states with a wrong (nonrelativistic) dispersion relation (see e.g. [85]).

3 Wavelet scaling of lattice scalar fields

3.1 Scalar fields on lattice

3.1.1 First and second quantization

The Weyl algebra is defined by the canonical commutation relations (a general reference is [11]). Let \mathfrak{h} be a Hilbert space (which will be called the **one-particle space**), $\langle \cdot, \cdot \rangle$ its scalar product and $\sigma(\cdot, \cdot) = \Im \langle \cdot, \cdot \rangle$, where \Im denotes the imaginary part, the canonical non degenerate symplectic form. A C^* -algebra $\mathcal{W}(\mathfrak{h})$ is said to be the **Weyl algebra associated to** (\mathfrak{h}, σ) if $\mathcal{W}(\mathfrak{h})$ is generated by elements $\{W(\xi)\}_{\xi \in \mathfrak{h}}$ with the following commutation relations:

$$W(\xi)W(\eta) = e^{-\frac{i}{2}\sigma(\xi, \eta)}W(\xi + \eta), \quad \xi, \eta \in \mathfrak{h}. \quad (3.1)$$

Such a C^* -algebra is unique (up to isomorphism) and simple.

A fundamental representation of $\mathcal{W}(\mathfrak{h})$ is constructed on the **bosonic Fock space over** \mathfrak{h} : $\mathfrak{F}_+(\mathfrak{h}) = \mathbb{C}\Omega \oplus \bigoplus_{n \in \mathbb{N}_0} \mathfrak{h}^{\otimes_s n}$, where $\mathfrak{h}^{\otimes_s n}$ is the symmetric n -fold tensor product of the one-particle space \mathfrak{h} and Ω is called the **Fock vacuum**. On $\mathfrak{F}_+(\mathfrak{h})$, the actions of the Weyl

operators $W(\xi)$ are uniquely given by

$$W(\xi)\Omega = e^{-\frac{1}{2}\|\xi\|^2} e^\xi, \quad \text{where } \xi \in \mathfrak{h} \text{ and } e^\xi = 1 \oplus \xi \oplus \frac{1}{2!}\xi^{\otimes 2} \oplus \dots \oplus \frac{1}{n!}\xi^{\otimes n} \oplus \dots$$

and by the canonical commutation relations (3.1): This is the GNS representation of the algebra $\mathcal{W}(\mathfrak{h})$ with respect to the state $\omega(W(\xi)) = \langle \Omega, W(\xi)\Omega \rangle = e^{-\frac{1}{2}\|\xi\|^2}$.

We use the following repeatedly, and hence state it as a proposition.

Proposition 3.1. *If we have a symplectic map R from \mathfrak{h}_1 into \mathfrak{h}_2 , we have a natural injective $*$ -homomorphism from $\mathcal{W}(\mathfrak{h}_1)$ to $\mathcal{W}(\mathfrak{h}_2)$, which maps $W(\xi)$ to $W(R(\xi))$, which is a $*$ -isomorphism if R is bijective.*

Note that, even if $\mathfrak{h}_1 \subset \mathfrak{h}_2$, we may take a symplectic map R which is different from the inclusion map. Indeed, such different embedding correspond to different scaling maps, see Section 3.3.

3.1.2 Scalar fields on different scales

As in Section 3.1.1, to construct the Weyl algebra, one has firstly to define the one-particle Hilbert space. Let us consider an initial lattice

$$\Lambda_{\varepsilon,r} = \varepsilon\{-r, \dots, 0, \dots, r-1\}^d \subset \mathbb{R}^d$$

with scale parameters³ $\varepsilon > 0$, $r \in \mathbb{N}$. We think of $\Lambda_{\varepsilon,r}$ as a discretization of the cube $[-L, L]^d = \mathbb{T}_L^d$ with periodic boundary conditions ($r \equiv -r$) which fixes the product of scale parameters $\varepsilon r = L > 0$. From $\Lambda_{\varepsilon,r}$ we generate a sequence of lattices $\Lambda_N = \Lambda_{\varepsilon_N, r_N}$ with $\varepsilon_N = 2^{-N}\varepsilon$, $r_N = 2^N r$ for $N \in \mathbb{N}_0$. This way all lattices have the same volume $\text{vol}(\Lambda_N) = (2\varepsilon_N r_N)^d = (2L)^d$. In the following, we also need the dual lattice

$$\Gamma_{\varepsilon,r} = \frac{\pi}{\varepsilon r}\{-r, \dots, 0, \dots, r-1\}$$

and its scaled versions $\Gamma_N = \Gamma_{\varepsilon_N, r_N}$, with the scaling parameter defined above.

We introduce two $2^N r$ -dimensional Hilbert spaces associated with the lattices and their dual⁴

$$\mathfrak{h}_{N,L} = \ell^2(\Lambda_N), \quad \langle \xi, \eta \rangle_{N,L} = \sum_{x \in \Lambda_N} \bar{\xi}(x)\eta(x), \quad (3.2)$$

$$\hat{\mathfrak{h}}_{N,L} = L^2(\Gamma_N, (2r_N)^{-d} \mu_{\Gamma_N}), \quad \langle \hat{\xi}, \hat{\eta} \rangle_{N,L} = (2r_N)^{-d} \sum_{k \in \Gamma_N} \bar{\hat{\xi}}(k)\hat{\eta}(k), \quad (3.3)$$

where the counting measure μ_{Γ_N} acquires a factor $(2r_N)^{-d}$ because of the normalization $(2r_N)^{-d} \sum_{k \in \Gamma_N} e^{ik \cdot (x-x')} = \delta_{x,x'}$, $x, x' \in \Lambda_N$.

These two Hilbert spaces are identified via the discrete Fourier transform:

$$\begin{aligned} \mathcal{F}_N[\xi](k) &= \hat{\xi}(k) = \sum_{x \in \Lambda_N} \xi(x)e^{-ik \cdot x}, \\ \mathcal{F}_N^{-1}[\hat{\xi}](x) &= \check{\xi}(x) = (2r_N)^{-d} \sum_{k \in \Gamma_N} \hat{\xi}(k)e^{ik \cdot x} = \xi(x). \end{aligned} \quad (3.4)$$

³The scale parameters ε, r will be fixed more explicitly below.

⁴We emphasize the dependence on the lattice size L only on Hilbert spaces and algebras, although most of other objects depend implicitly on L .

The kinematical scalar field lattice systems are given in terms of the one-particle spaces $\mathfrak{h}_{N,L}$, cp. [4, 11]:

$$\mathcal{W}(\mathfrak{h}_{N,L}) = \mathcal{W}_{N,L}, \quad \mathcal{H}_{N,L} = \mathfrak{F}_+(\mathfrak{h}_{N,L}) \cong \bigotimes_{x \in \Lambda_N} \mathcal{H}_x, \quad (3.5)$$

where $\mathcal{H}_x = L^2(\mathbb{R})$, and $\mathcal{W}(\mathfrak{h}_{N,L})$ is the Weyl algebra,

$$W_N(\xi)W_N(\eta) = e^{-\frac{i}{2}\sigma_{N,L}(\xi,\eta)}W_N(\xi + \eta), \quad \xi, \eta \in \mathfrak{h}_{N,L}, \quad (3.6)$$

with respect to the standard symplectic form, $\sigma_{N,L} = \mathfrak{S}\langle \cdot, \cdot \rangle_{\mathfrak{h}_{N,L}}$. The decomposition into real (Langragian) subspaces is facilitated by

$$\xi = \varepsilon_N^{\frac{d+1}{2}} q_\xi + i\varepsilon_N^{\frac{d-1}{2}} p_\xi \text{ for } \xi \in \mathfrak{h}_{N,L} \quad (3.7)$$

with real-valued $q_\xi, p_\xi \in \mathfrak{h}_{N,L}$. We denote by q, p generic real-valued elements in $\mathfrak{h}_{N,L}$. The finite volume field $\Phi_N(q)$ and momentum $\Pi_N(p)$ are the generators of the one-parameter groups $W_N(tq), W_N(tp)$, respectively, and in terms of (q, p) the symplectic form reads

$$\sigma_{N,L}((q_\xi, p_\xi), (q_\eta, p_\eta)) := \sigma_{N,L}(\xi, \eta) = \varepsilon_N^d \sum_{x \in \Lambda_N} q_\xi(x)p_\eta(x) - p_\xi(x)q_\eta(x). \quad (3.8)$$

For the Lagrangian decomposition of $\hat{\mathfrak{h}}_{N,L}$, we choose the normalization:

$$\hat{q} = \varepsilon_N^{\frac{d}{2}} \mathcal{F}_N[q], \quad \hat{p} = \varepsilon_N^{\frac{d}{2}} \mathcal{F}_N[p]. \quad (3.9)$$

The algebra $\mathcal{W}_{N,L}$ is also naturally equipped with a *-automorphic action $\eta_L^{(N)} : \Lambda_N \curvearrowright \mathcal{W}_{N,L}$ of lattice translations, induced via Prop. 3.1 by the action $\tau_L^{(N)} : \Lambda_N \curvearrowright \mathfrak{h}_{N,L}$ of translations on the one-particle space, defined by the symplectic maps $(\tau_{L|a}^{(N)}\xi)(x) := \xi(x-a)$, $a, x \in \Lambda_N$, $\xi \in \mathfrak{h}_{N,L}$, where the difference $x - a$ is to be intended mod $2L\mathbb{Z}^d$.

3.1.3 Scaling maps from one-particle spaces

A renormalization group for lattice scalar fields (see Section 3.1.2) is obtained from a symplectic map $R_{N'}^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_{N',L}$ between one-particle spaces: it induces via Prop. 3.1 an injective *-homomorphism $\alpha_{N'}^N : \mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N',L}$ such that

$$\alpha_{N'}^N(W_N(\xi_N)) = W_{N'}(R_{N'}^N(\xi_N)). \quad (3.10)$$

We thus obtain a C^* -inductive system of lattice Weyl algebras,

$$\mathcal{W}_{0,L} \rightarrow \cdots \rightarrow \mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N+1,L} \rightarrow \cdots \rightarrow \mathcal{W}_{N',L} \rightarrow \cdots. \quad (3.11)$$

Although there is a natural inclusion $\mathfrak{h}_{N,L} \subset \mathfrak{h}_{N+1,L}$, we *do not* take it as R_{N+1}^N , because this would mean (cf. Section 3.2) that we identify an operator on a lattice vertex with an operator localized in one point in the continuum, which does not exist. Instead, we consider maps associated with various wavelets.

As Weyl algebras are simple, thus *-homomorphisms are injective, and the inductive limit of a directed system of simple C^* -algebras is again simple.

3.2 An instructive example: The block-spin method

As an illustration of the general scheme introduced above, we define a version of the block-spin renormalization group for lattice scalar fields known from spin systems (e.g. the Ising model), see [28].

Definition 3.2. *The block-spin renormalization group $\{\alpha_{N'}^N\}_{N < N'}$ is the inductive family of *-homomorphisms defined by the block-spin scaling map between one-particle Hilbert spaces:*

$$R_{N+1}^N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{h}_{N+1,L},$$

where

$$R_{N+1}^N(q, p)(x') = \sum_{x \in \Lambda_N} (q, p)(x) \chi_{[0, \varepsilon_N]^d}(x' - x), \quad N \in \mathbb{N}_0, \quad (3.12)$$

(note that q and p in $\mathfrak{h}_{N,L}$ are scaled as (3.7), that compensate the volume of the cube) namely, the function with support in x is mapped to the step function supported in the (discrete) cube with “lower left” corner x . Other scaling maps are defined by composition, so that the semigroup property is automatic:

$$R_{N'}^N = R_{N'}^{N'-1} \circ R_{N'-1}^{N'-2} \circ \dots \circ R_{N+1}^N, \quad N < N'.$$

We note that R_{N+1}^N is symplectic, because the inner product is preserved:

$$\sigma_{N+1,L}(R_{N+1}^N(q, p), R_{N+1}^N(q', p')) = \sigma_{N,L}((q, p), (q', p')),$$

Intuitively, (3.12) encodes the idea that the field and its momentum spatially localized at x at scale N result from averaging⁵ over points x' close by at scale $N + 1$. At the level of fields, this yields:

$$\alpha_{N+1}^N(\Phi_N(x)) = 2^{-d} \sum_{x' \in \Lambda_{N+1}} \chi_{[0, \varepsilon_N]^d}(x' - x) \Phi_{N+1}(x'), \quad (3.13)$$

$$\alpha_{N+1}^N(\Pi_N(x)) = 2^{-d} \sum_{x' \in \Lambda_{N+1}} \chi_{[0, \varepsilon_N]^d}(x' - x) \Pi_{N+1}(x'). \quad (3.14)$$

Formally, the block-spin scaling map encodes the relation between block-averaged (continuum) fields and momenta at scales $N \in \mathbb{N}_0$, i.e. we think of the lattice fields and momenta as integrated against characteristic functions of lattice cubes with Φ, Π the continuum field and the continuum momenta,

$$\Phi_N(x) = \varepsilon_N^{-d} \int_{\mathbb{T}_L^d} d^d x' \chi_{[0, \varepsilon_N]^d}(x' - x) \Phi(x') = \varepsilon_N^{-d} \Phi(\chi_{x+[0, \varepsilon_N]^d}) \text{ (formal)}, \quad (3.15)$$

$$\Pi_N(x) = \varepsilon_N^{-d} \int_{\mathbb{T}_L^d} d^d x' \chi_{[0, \varepsilon_N]^d}(x' - x) \Pi(x') = \varepsilon_N^{-d} \Pi(\chi_{x+[0, \varepsilon_N]^d}) \text{ (formal)}, \quad (3.16)$$

although Π is ill-defined for nondifferentiable functions.

Yet, the idea of embedding lattice fields into the continuum field can be justified using wavelets. Indeed, if the problem is the non-differentiability of step functions, we would only have to identify lattice fields with continuum fields smeared over more regular functions, as we see below.

⁵Since we are working with the algebra of field operators, the block-spin transformation results in a refining operation $\mathfrak{A}_N = \mathcal{W}_{N,L} \rightarrow \mathfrak{A}_{N+1} = \mathcal{W}_{N+1,L}$ (2.4) in contrast with the familiar coarse-graining operation on states, respectively density matrices (2.2).

3.3 A generalization: The wavelet method

3.3.1 Scaling map from scaling function.

The block-spin scaling map is a special case of wavelet scaling map which we define here. The scaling map is induced by the maps between lattice one-particle spaces:

$$\mathfrak{h}_{0,L} \cdots \xrightarrow{R_N^{N-1}} \mathfrak{h}_{N,L} \xrightarrow{R_{N+1}^N} \mathfrak{h}_{N+1,L} \xrightarrow{R_{N+2}^{N+1}} \cdots \xrightarrow{R_{N'+1}^{N'-1}} \mathfrak{h}_{N',L} \xrightarrow{R_{N'+2}^{N'}} \cdots, \quad (3.17)$$

As we hope to embed lattice fields into the continuum field, there should be the corresponding spaces in the continuum one-particle space:

$$V_0 \subset \cdots \subset V_N \subset V_{N+1} \subset \cdots \subset L^2(\mathbb{T}_L^d). \quad (3.18)$$

In the case of the block-spin renormalization, V_N is spanned by the step functions $\chi_{[0,\varepsilon_n)^d}(\cdot - x)$, and this is naturally included in V_{N+1} . Furthermore, such a step function is a linear combination of finer step functions with a fixed set of coefficients (see Section 3.3.3 for details of this case). By generalizing these properties, we are led to consider wavelets. A general reference is [23] (for periodic wavelet bases, see [23, Section 10.7]).

We start with a $2r$ -periodic **scaling function** $\phi \in L^2(\mathbb{T}_r^d)$, and define the rescaled scaling function $\phi^{(\varepsilon)}(x) := \varepsilon^{-d/2} \phi(\varepsilon^{-1}x)$, $x \in \mathbb{T}_L^d$ (recall that $L = \varepsilon r$). The characteristic properties of ϕ translate into the following ones⁶ for the function $\phi^{(\varepsilon)}$.

- $\{\phi^{(\varepsilon)}(\cdot - k)\}_{k \in \Lambda_{\varepsilon,r}}$ is an orthonormal system.
- With the (normalized) rescaled functions $\phi_{N,k}^{(\varepsilon)}(x) = 2^{\frac{Nd}{2}} \phi^{(\varepsilon)}(2^N x - k)$, $N \in \mathbb{N}_0$, $k \in \Lambda_{\varepsilon,r_N} = \{-\varepsilon r_N, \dots, \varepsilon(r_N - 1)\}^d \subset \varepsilon \mathbb{Z}^d$, it holds that

$$\phi^{(\varepsilon)}(x) = \sum_{n \in \Lambda_{1,r_1}} h_n \phi_{1,\varepsilon n}^{(\varepsilon)}(x), \quad (3.19)$$

with $h_n \in \mathbb{R}$.

- $\{\phi_{N,k}^{(\varepsilon)}\}_{N \in \mathbb{N}_0, k \in \Lambda_{\varepsilon,r_N}}$ span $L^2(\mathbb{T}_L^d)$.
- $\phi^{(\varepsilon)}$ is normalized in such a way that

$$\int_{\mathbb{T}_L^d} \phi^{(\varepsilon)}(x) dx = \varepsilon^{d/2}. \quad (3.20)$$

We discuss concrete examples of scaling functions in the next sections. In case the scaling function ϕ is chosen compactly supported, there is $r_{\max} > 0$ such that $h_n \neq 0$ only for $\|n\|_{\infty} \leq r_{\max}$, where $\|n\|_{\infty}$ is the largest absolute value of the components of $n \in \Lambda_{1,r_1} \subset \mathbb{Z}^d$, because these are the expansion coefficients of $\phi^{(\varepsilon)}$ in the basis $\{\phi_{1,\varepsilon n}^{(\varepsilon)}\}$, i.e. $\{h_n\}_{\|n\|_{\infty} \leq r_{\max}}$ yields a finite impulse response filter scheme, and is called a low-pass filter [23].

⁶Here the convention is that n belongs to a unit lattice (with lattice spacing 1) while k belongs to a scaled lattice.

A (half-sided) multiresolution analysis (MRA)⁷ is associated to $\phi^{(\varepsilon)}$, i.e. the sequence of subspaces V_N spanned by $\{\phi_{N,k}^{(\varepsilon)}\}_{k \in \Lambda_{\varepsilon,r_N}}$, $N \in \mathbb{N}_0$, satisfying (3.18) with the properties:

$$\overline{\bigcup_{N \in \mathbb{N}_0} V_N} = L^2(\mathbb{T}_L^d), \quad (3.21)$$

$$\bigcap_{N \in \mathbb{N}_0} V_N = V_0, \quad (3.22)$$

$$f \in V_0 \Leftrightarrow f(2^N \cdot) \in V_N, \quad (3.23)$$

$$f \in V_0 \Leftrightarrow f(\cdot - k) \in V_0 \quad \forall k \in \varepsilon \mathbb{Z}^d. \quad (3.24)$$

For a fixed N , $\{\phi_{N,k}^{(\varepsilon)}\}_{k \in \Lambda_{\varepsilon,r_N}}$ play the role of approximate δ -functions. We will also use the notation $\phi_x^{(\varepsilon_N)} := \phi_{N,2^N x}^{(\varepsilon)}$, $x \in \Lambda_N (= \Lambda_{2^{-N}\varepsilon, 2^N r})$, $N \in \mathbb{N}_0$. Correspondingly, we denote the standard basis of $\ell^2(\Lambda_N)$ by $\{\delta_x^{(N)}\}_{x \in \Lambda_N}$. At this point, we demand $\log_2 r \in \mathbb{N}_0$ as otherwise the completeness of the restricted half-sided MRA (3.18) to $L^2(\mathbb{T}_L^d)$ is problematic [23]. Having in mind the identification $\delta_x^{(0)} \sim \phi_x^{(\varepsilon)}$, $x \in \Lambda_{\varepsilon,r} = \Lambda_0$, we make the analogue of Definition 3.2.

Definition 3.3. *Given a scaling function $\phi^{(\varepsilon)}$ with the scaling coefficients (3.19), the wavelet renormalization group $\{\alpha_{N'}^N\}_{N < N'}$ is the inductive family of *-homomorphisms defined by the wavelet scaling map between one-particle Hilbert spaces:*

$$R_{N+1}^N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{h}_{N+1,L},$$

$$R_{N+1}^N(q,p)(x) = 2^{\frac{d}{2}} \sum_{x' \in \Lambda_N} (q,p)(x') \sum_{n \in \mathbb{Z}^d, \|n\|_{\infty} \leq r_{\max}} h_n \delta_{x'+n\varepsilon_{N+1}}^{(N+1)}(x), \quad (3.25)$$

where the index of $\delta_{x'+n\varepsilon_{N+1}}^{(N+1)} \in \ell^2(\Lambda_{N+1})$ is interpreted $\text{mod } 2r_{N+1}(\varepsilon_{N+1}\mathbb{Z}^d)$, according to the periodization convention above, and

$$R_{N'}^N = R_{N'}^{N'-1} \circ R_{N'-1}^{N'-2} \circ \dots \circ R_{N+1}^N, \quad N < N'.$$

The numerical coefficients in (3.25) are motivated by the formal relations between fields and momenta at successive scales implied by (3.19), cp. (3.13) and (3.14). In the following, to simplify the notation, we will indicate sums as the second one in (3.25) simply as sums over $n \in \mathbb{Z}^d$, with the convention that $h_n = 0$ for $\|n\|_{\infty} > r_{\max}$.

Thanks to the property (3.19), we obtain a family of symplectic maps.

Lemma 3.4. *The scaling maps $R_{N'}^N$, $N < N'$, associated with a scaling function ϕ are symplectic, i.e.*

$$\sigma_{N',L}(R_{N'}^N(q,p), R_{N'}^N(q',p')) = \sigma_{N,L}((q,p), (q',p')).$$

⁷In the non-periodic setting this sequence is two-sided infinite, i.e. indexed by \mathbb{Z} .

Proof. A direct computation for R_{N+1}^N with $N \in \mathbb{N}_0$ arbitrary yields:

$$\begin{aligned}
& \varepsilon_{N+1}^d \sum_{x' \in \Lambda_{N+1}} R_{N+1}^N(p)(x') R_{N+1}^N(q')(x') \\
&= \frac{\varepsilon_N^d}{2^d} \sum_{x' \in \Lambda_{N+1}} \left(2^{\frac{d}{2}} \sum_{x \in \Lambda_N} p(x) \sum_{n \in \mathbb{Z}^d} h_n \delta_{x+n\varepsilon_{N+1}}^{(N+1)}(x') \right) \left(2^{\frac{d}{2}} \sum_{y \in \Lambda_N} q'(y) \sum_{m \in \mathbb{Z}^d} h_m \delta_{y+m\varepsilon_{N+1}}^{(N+1)}(x') \right) \\
&= \varepsilon_N^d \sum_{x, y \in \Lambda_N} p(x) q'(y) \sum_{n, m \in \mathbb{Z}^d} h_n h_m \delta_{n-m, 2\varepsilon_N^{-1}(y-x)} \\
&= \varepsilon_N^d \sum_{x, y \in \Lambda_N} p(x) q'(y) \underbrace{\sum_{n \in \mathbb{Z}_r^d} h_n h_{n+2\varepsilon_N^{-1}(x-y)}}_{=\delta_{x,y}} \\
&= \varepsilon_N^d \sum_{x \in \Lambda_N} p(x) q'(x),
\end{aligned}$$

where we used the fact that $2\varepsilon_N^{-1}(x-y) \in 2\mathbb{Z}^d$ for $x, y \in \Lambda_N$ and the orthogonality relation $\sum_{n \in \mathbb{Z}^d} h_n h_{n+2m} = \delta_{m,0}$ which follows from the scaling equation (3.19) and the orthonormality of $\{\phi_{0,k}^{(\varepsilon)}\}_{k \in \Lambda_{\varepsilon,r}}$. The statement is then obtained by (3.8). \square

Therefore, according to the discussion in Section 3.1.3, we denote by $\mathcal{W}_{\infty,L}$ the inductive limit C^* -algebra obtained from the inductive system defined by (3.10), (3.11). Moreover, it is easy to check that for the one-particle lattice translations $\tau_{L|a}^{(N)}$, $a \in \Lambda_N$, there holds $\tau_{L|a}^{(N')} \circ R_{N'}^N = R_{N'}^N \circ \tau_{L|a}^{(N)}$, which of course entails, for the lattice translation automorphisms $\eta_{L|a}^{(N)}$ of $\mathcal{W}_{N,L}$,

$$\eta_{L|a}^{(N')} \circ \alpha_{N'}^N = \alpha_{N'}^N \circ \eta_{L|a}^{(N)}, \quad a \in \Lambda_N, N' > N. \quad (3.26)$$

This, together with the inductive limit uniqueness, implies the existence of an automorphic action of the dyadic translations on the inductive limit algebra as follows. Given $a \in \bigcup_N \Lambda_N$ and $N' \in \mathbb{N}$, since $\Lambda_N \subset \Lambda_{N+1}$ we can assume that $a \in \Lambda_N$ with $N \geq N'$, and we can define an injective $*$ -morphism $\beta_{\infty}^{N'} : \mathcal{W}_{N',L} \rightarrow \mathcal{W}_{\infty,L}$ by

$$\beta_{\infty}^{N'}(A) := \alpha_{\infty}^N(\eta_{L|a}^{(N)}(\alpha_{N'}^N(A))), \quad A \in \mathcal{W}_{N',L}.$$

Thanks to the intertwining property (3.26), $\beta_{\infty}^{N'}$ is independent of the chosen N such that $a \in \Lambda_N$, and one immediately checks that $\beta_{\infty}^{N'} \circ \alpha_{N'}^{N''} = \beta_{\infty}^{N''}$ for $N' \geq N''$. Therefore, by the uniqueness of the C^* -inductive limit, there exists a $*$ -automorphism $\eta_{L|a}^{(\infty)} : \mathcal{W}_{\infty,L} \rightarrow \mathcal{W}_{\infty,L}$ such that $\eta_{L|a}^{(\infty)} \circ \alpha_{\infty}^{N'} = \beta_{\infty}^{N'}$ for all $a \in \bigcup_N \Lambda_N$ and $N' \in \mathbb{N}$, i.e.,

$$\eta_{L|a}^{(\infty)}(\alpha_{\infty}^{N'}(A)) = \alpha_{\infty}^N(\eta_{L|a}^{(N)}(\alpha_{N'}^N(A))), \quad A \in \mathcal{W}_{N',L}, a \in \Lambda_N, N \geq N', \quad (3.27)$$

and $\eta_L^{(\infty)} : \bigcup_N \Lambda_N \curvearrowright \mathcal{W}_{\infty,L}$ defines the required action of the dyadic translation group.

Invoking the discrete Fourier transform (3.4), we obtain the important identity in momentum space, i.e. with respect to the inclusion of dual lattices $\Gamma_N \subset \Gamma_{N+1}$:

$$R_{N+1}^N(\hat{q}_N, \hat{p}_N) = 2^{\frac{d}{2}} m_0(\varepsilon_{N+1} \cdot)(\hat{q}_{N+1}, \hat{p}_{N+1}), \quad (3.28)$$

where $m_0(\varepsilon_{N+1}k) = 2^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n e^{-i\varepsilon_{N+1}nk}$ (the discrete Fourier transform of the coefficients h_n) and $\hat{q}_{N+1}, \hat{p}_{N+1}$ are the periodic extension of \hat{q}_N, \hat{p}_N from Γ_N to Γ_{N+1} . From (3.28), we easily compute the iterated maps,

$$R_{N'}^N(\hat{q}_N, \hat{p}_N) = 2^{\frac{d(N'-N)}{2}} \prod_{n=1}^{N'-N} m_0(\varepsilon_{N+n} \cdot)(\hat{q}_{N'}, \hat{p}_{N'}). \quad (3.29)$$

This expression already indicates the special role of the scaling function ϕ in the limit $N' \rightarrow \infty$ (if r_{\max} is finite) because [23, (6.2.2)]:

$$\lim_{N' \rightarrow \infty} \prod_{n=1}^{N'-N} m_0(\varepsilon_{N+n} \cdot) = \hat{\phi}(\varepsilon_N \cdot) = \varepsilon^{-d/2} \hat{\phi}(\varepsilon)(2^{-N} \cdot) = \varepsilon_N^{-d/2} \hat{\phi}_0(\varepsilon_N) \quad (3.30)$$

pointwise, which results from the (continuum) Fourier transform (normalized as in (4.12)) of the scaling equation (3.19):

$$\hat{\phi}(\varepsilon)(2^{-N}k) = m_0(\varepsilon_{N+1}k) \hat{\phi}(\varepsilon)(2^{-(N+1)}k), \quad k \in \frac{\pi}{L} \mathbb{Z}^d, \quad (3.31)$$

together with the normalization (3.20). Thus, apart from an infinite field-strength renormalization accounting for the rescaling of the symplectic structure $\sigma_{N',L}$ with N' , the inductive limit map, $R_{\infty}^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_{\infty,L} = \varinjlim_N \mathfrak{h}_{N,L}$, is given by multiplication with the Fourier transform $\varepsilon_N^{-\frac{d}{2}} \hat{\phi}_0(\varepsilon_N)$ in momentum space.

3.3.2 Wavelet bases.

Before turning to a detailed description of the wavelet method, let us also comment on the use of the wavelet basis of $L^2(\mathbb{T}_L^d)$ constructed from the scaling function (3.37). In general, such a basis is obtained by rewriting the MRA (3.18) as a direct sum decomposition:

$$V_0 \oplus V'_0 \oplus \dots \oplus V'_N \oplus V'_{N+1} \oplus \dots = L^2(\mathbb{T}_L^d). \quad (3.32)$$

This is achieved by finding an orthogonal decomposition of the orthogonal complement of V_0 inside V_1 :

$$V_1 = V_0 \oplus V'_0 = V_0 \oplus \bigoplus_{m=1}^{2^d-1} V_0^{\prime,m}, \quad (3.33)$$

such that there are distinguished orthonormal functions – the **wavelets**, $\{\psi_0^{(\varepsilon)}\}_{m=1}^{2^d-1}$, which together with their integer translates span the spaces $V_0^{\prime,m}$. The spaces $V_N^{\prime,m}$ are constructed by rescaling the wavelets, thus, these spaces inherit the scaling properties of the V_N .

As the use of wavelets compared to that of the scaling function turns out to be important for the characterization of the inductive-limit algebra $\mathcal{W}_{\infty,L}$ and the scaling limit $\omega_{\infty}^{(\infty)} = \omega_{L,\infty}^{(\infty)}$ (2.11) (see Section 4), we also provide some details on the construction of wavelets.

Let us discuss wavelets in the unit lattice, $\varepsilon = 1$ for the case $d = 1$. Higher dimensional wavelets can be obtained by tensor product, cf. [23].

Given the scaling function, $\phi_0^{(\varepsilon=1)} = \phi$, the wavelet $\psi_0^{(\varepsilon=1)} = \psi$ can be constructed by the formula:

$$\psi(x) = \sum_{n \in \mathbb{Z}} g_n \phi_{1,n}(x),$$

where the wavelet coefficients $\{g_n\}_{n \in \mathbb{Z}}$ (also called a high-pass filter) are obtained explicitly in terms of the expansion coefficients $\{h_n\}_{n \in \mathbb{Z}}$ of the scaling equation (3.39). Albeit, the choice of wavelet coefficients is not unique, convenient options include:

$$g_n = (-1)^n \overline{h_{1-n}}, \quad \text{or} \quad g_n = (-1)^n h_{1-n+2M}, \quad \forall n \in \mathbb{Z}, \quad (3.34)$$

for some $M \in \mathbb{Z}$, and we take one of them. By construction, the set of functions $\{\phi\} \cup \{\psi_{N,k}\}_{N \in \mathbb{N}_0, k \in \{-2^N r, \dots, 2^N r - 1\}}$, where $\psi_{N,k}(x) = 2^{\frac{Nd}{2}} \psi(2^N x - k)$, forms an orthonormal basis of $L^2(\mathbb{T}_{L=r})$. The orthogonal decomposition $V_{N+1} = V_N \oplus V'_N$ analogous to (3.33) is captured by an important projection identity involving the wavelets $\psi_{N,k}$ and scaling functions $\phi_{N,k}$:

$$\begin{aligned} \text{Proj}_{V_{N+1}}(f) &= \sum_{k=-2^{N+1}r}^{2^{N+1}r-1} \langle \phi_{N+1,k}, f \rangle_{L^2} \phi_{N+1,k} \\ &= \text{Proj}_{V_N}(f) + \sum_{k=-2^N r}^{2^N r-1} \langle \psi_{N,k}, f \rangle_{L^2} \psi_{N,k} \\ &= \sum_{k=-2^N r}^{2^N r-1} \langle \phi_{N,k}, f \rangle_{L^2} \phi_{N,k} + \sum_{k=-2^N r}^{2^N r-1} \langle \psi_{N,k}, f \rangle_{L^2} \psi_{N,k}, \quad f \in L^2(\mathbb{T}_L). \end{aligned} \quad (3.35)$$

Iterating and using $\lim_{N \rightarrow \infty} \text{Proj}_{V_N} = \text{id}_{L^2(\mathbb{T}_L)}$, we have an expansion:

$$f = \sum_{k=-r}^{r-1} \langle \phi_{0,k}, f \rangle_{L^2} \phi_{0,k} + \sum_{N \in \mathbb{N}_0} \sum_{k=-2^N r}^{2^N r-1} \langle \psi_{N,k}, f \rangle_{L^2} \psi_{N,k}. \quad (3.36)$$

3.3.3 Block-spin renormalization in terms of orthogonal Haar wavelets

The above interpretation (3.15) and (3.16) of lattice fields as continuum field smeared with characteristic functions, $\{\varepsilon_N^{-d} \chi_{x+[0, \varepsilon_N]^d}\}_{x \in \Lambda_N}$, can be understood as a special instance of the general scheme of wavelet scaling discussed above. We take the characteristic function $\chi_{[0, \varepsilon]^d}$ as the scaling function underlying the construction of the (periodic) Haar wavelets basis of $L^2(\mathbb{T}_L^d)$ with scaling parameters $\varepsilon r = L$ [4, 58, 19, 23]. More precisely, the function,

$$\phi_x^{(\varepsilon)}(y) = \varepsilon^{-\frac{d}{2}} \chi_{[0, \varepsilon]^d}(y - x) = \varepsilon^{-\frac{d}{2}} \chi_{[0, 1]^d}(\varepsilon^{-1}(y - x)), \quad (3.37)$$

has an associated (half-sided) multiresolution analysis (MRA), i.e. there is sequences of subspaces of step functions $\{V_N\}_{N \in \mathbb{N}_0}$ satisfying (3.21)-(3.24). In this case, we have $r_{\max} = 1$.

By dimensionality, we can identify the real subspace spanned by this basis $\{\phi_{N,k}^{(\varepsilon)}\}$ with the Lagrangian subspaces of $\mathfrak{h}_{N,L}$, cp. Section 3.1.2. Moreover, Definition 3.2 of the block-spin scaling map can be obtained from the expansion of $\phi_0^{(\varepsilon)}$ into the functions $\{\phi_{1,k}^{(\varepsilon)}\}$, which is equivalent to the inclusion $V_0 \subset V_1$ and similarly, an embedding $\mathfrak{h}_{0,L} \rightarrow \mathfrak{h}_{1,L}$ is given by writing a step function as a linear combination of finer step functions:

$$\phi_0^{(\varepsilon)}(x) = \sum_{k \in \mathbb{N}_0^d, \|k\|_\infty \leq r_{\max}} 2^{-\frac{d}{2}} \phi_{1, \varepsilon k}^{(\varepsilon)}(x). \quad (3.38)$$

Now, (3.38) can be interpreted as the well-known scaling equation,

$$\phi_0^{(\varepsilon)}(x) = \sum_{n \in \mathbb{Z}^d} h_n \phi_{1, \varepsilon n}^{(\varepsilon)}(x), \quad (3.39)$$

of the theory of wavelets associated with the scaling function (3.37) (and the MRA (3.18)). The Haar wavelet basis in $L^2([0, 1])$ ($d = 1$) is usually given as:

$$\psi_{0,0}(x) = (-1)\phi(2x - 1) + (+1)\phi(2x) = \begin{cases} 1 & x \in [0, \frac{1}{2}) \\ -1 & x \in [\frac{1}{2}, 1), \end{cases}$$

$$\psi_{N,k}(x) = 2^{\frac{N}{2}}\psi_{0,0}(2^N x - k), \quad N \in \mathbb{N}_0, 0 \leq k \leq 2^N - 1.$$

The relation between fields and momenta at successive scales (3.13) and (3.14) in terms of the scaling function $\phi_0^{(\varepsilon)}$ can be reformulated as follows:

$$\Phi_N(x) = \varepsilon_N^{-\frac{d}{2}}\Phi(\phi_x^{(\varepsilon_N)}) = 2^{-\frac{d}{2}}\varepsilon_{N+1}^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n \Phi(\phi_{1,2x+\varepsilon_N n}^{(\varepsilon_N)}) \quad (3.40)$$

$$= 2^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n \varepsilon_{N+1}^{-\frac{d}{2}} \Phi(\phi_{x+\varepsilon_{N+1} n}^{(\varepsilon_{N+1})}) = 2^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n \Phi_{N+1}(x + \varepsilon_{N+1} n),$$

$$\Pi_N(x) = 2^{-\frac{d}{2}} \sum_{n \in \mathbb{Z}^d} h_n \Pi_{N+1}(x + \varepsilon_{N+1} n). \quad (3.41)$$

3.3.4 The Daubechies wavelets

In our renormalization scheme of Section 2.4, observables in the lattice field theories should be mapped to certain observables in the continuum theory. In order to obtain a continuum field as the scaling limit of lattice fields, we need to choose a scaling function ϕ which is both localized and sufficiently regular. This goal is achieved by the so-called Daubechies wavelets [23, Chapter 6].

This is a family of scaling functions $\{{}_K\phi\}_{K \in \mathbb{N}}$ with various support properties and regularity, although no closed expression for them is known. The scaling function ${}_K\phi$ satisfies the (3.39) with $\{h_n\}$ with $h_n = 0$ for $n \geq 2K$ [23, Table 6.1], is supported in $[0, 2K - 1]$ and belongs to the class $C^{\alpha-\epsilon}$ for arbitrary $\epsilon > 0$, where the dependence of α on K is given by [23, Table in P.226]. In particular, with $K = 2$, $\alpha \cong 0.339$, and this suffices for our purpose to construct both the field and momentum operators in the continuum.

In the following, we take $\phi = {}_K\phi$ and define R_{N+1}^N as in Section (3.25).

4 The continuum limit of the free vacua

On the field algebras $\{\mathcal{W}_{N,L}\}$ on the lattices with the family of scaling maps $\{\alpha_{N'}^N\}_{N < N'}$ based on the Daubechies wavelets, we construct a family of initial states $\{\omega_{L,0}^{(N)}\}_{N \in \mathbb{N}_0}$, consisting of the initial data of Wilson's triangle of renormalization (see Figure 2). We restrict our attention to free fields, i.e. we choose as initial states a family of ground states of the free lattice Hamiltonian $H_{L,0}^{(N)}$.

As $\{\mathcal{W}_{N,L}\}$ can be interpreted as the time-zero algebra of the lattice field $\Phi_N(x), \Pi_N(x)$ and δ -functions on the lattice Λ_N are mapped to the scaling functions by (3.30), a simplest choice is to take the vacuum state in the continuum ω_L and to restrict it to the image of $\{\mathcal{W}_{N,L}\}$ by the map. It is straightforward to show that this yields a time-zero net of local algebras (by using the properties of the continuum free field). This is, however, available just because we know explicitly what the continuum state should look like. In a constructive program, one should take a natural Hamiltonian at each scale, and consider the sequence of the ground states. Below we show that this indeed gives the continuum free field state as well in the scaling limit, and hence gives rise to a continuum net.

4.1 States on lattice and continuum fields

4.1.1 Ground states of lattice free fields

The (self-adjoint) free lattice Hamiltonian $H_{L,0}^{(N)}$ of (unrenormalized) mass $\mu_N > \sqrt{2d}$ is defined on a dense domain $\mathcal{D}_{N,L} \subset \mathcal{H}_{N,L} = \mathfrak{F}_+(\mathfrak{h}_{N,L})$ by the expression [4, (1.8.17)] (up to a reparametrization) :

$$H_{L,0}^{(N)} = \frac{1}{2}\varepsilon_N^d \left(\sum_{x \in \Lambda_N} \Pi_{N|x}^2 + \mu_N^2 \varepsilon_N^{-2} \Phi_{N|x}^2 - 2 \sum_{\substack{x,y \in \Lambda_N \\ \text{adjacent}}} \varepsilon_N^{-2} \Phi_{N|x} \Phi_{N|y} \right). \quad (4.1)$$

To be precise, we define the domain $\mathcal{D}_{N,L} = \mathcal{D}_{\mu_N}$ depending explicitly on μ_N via the dispersion relation, $\gamma_{\mu_N}^2(k) = \varepsilon_N^{-2}(\mu_N^2 - 2d) + 2\varepsilon_N^{-2} \sum_{j=1}^d (1 - \cos(\varepsilon_N k_j))$, $k \in \Gamma_N$:

$$\mathcal{D}_{\mu_N} = \left\{ \Psi \in \mathcal{H}_{N,L} : \sum_{n=0}^{\infty} \left\| \sum_{j=1}^n \gamma_{\mu_N}((\cdot)_j) \Psi_n \right\|_{\ell^2(\Gamma_N)^{\otimes n}}^2 < \infty \right\}, \quad (4.2)$$

where $\gamma_{\mu_N}((\cdot)_j)$ denotes the multiplication operator on the j -th tensor component. The generators of the Weyl algebra $\mathcal{W}_{N,L}$ are related to the field, Φ_N , and momentum, Π_N , in the usual way (in the Fock representation on $\mathcal{H}_{N,L}$): $W_N(\xi) = e^{i(\Phi_N(q) + \Pi_N(p))}$ with $\xi = \varepsilon_N^{\frac{d+1}{2}} q + i\varepsilon_N^{\frac{d-1}{2}} p$ as in Section 3.1.2. The ground state (or lattice vacuum) Ω_{μ_N} of $H_{L,0}^{(N)}$ gives the following state on $\mathcal{W}_{N,L}$:

$$\omega_{\mu_N}(W_N(\xi)) = e^{-\frac{1}{4} \left(\left\| \gamma_{\mu_N}^{-1/2} \hat{q} \right\|_{\mathfrak{h}_{N,L}}^2 + \left\| \gamma_{\mu_N}^{1/2} \hat{p} \right\|_{\mathfrak{h}_{N,L}}^2 \right)}, \quad (4.3)$$

and the GNS construction applied to $\mathcal{W}_{N,L}$ with respect to ω_{μ_N} yields a representation which is unitarily equivalent to that on $\mathcal{H}_{N,L}$ such that Ω_{μ_N} is identified with the cyclic GNS vector.

Let us introduce the (N -dependent) physical mass m by $\mu_N^2 = \varepsilon_N^2 m^2 + 2d$. The Fock representation of mass m of the lattice scalar field $\mathcal{W}_{N,L}$ is determined by the Fock vacuum state⁸

$$\omega_{L,0}^{(N)}(W_N(\xi)) = e^{-\frac{1}{2} \left\| \hat{\xi}^{(m)} \right\|_{\mathfrak{h}_{N,L}}^2}, \quad \xi \in \mathfrak{h}_{N,L}, \quad (4.4)$$

where

$$\hat{\xi}^{(m)}(k) = \frac{i}{\sqrt{2}} \left(\gamma_m^{(N)}(k)^{-\frac{1}{2}} \hat{q}(-k) + i\gamma_m^{(N)}(k)^{\frac{1}{2}} \hat{p}(-k) \right)$$

with $\gamma_m^{(N)}(k)^2 = m^2 + 2\varepsilon_N^{-2} \sum_{j=1}^d (1 - \cos(\varepsilon_N k_j))$, $k \in \Gamma_N$. We actually have

$$\begin{aligned} \left\| \hat{\xi}^{(m)} \right\|_{\mathfrak{h}_{N,L}}^2 &= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_N} \left(\gamma_m^{(N)}(k)^{-1} \hat{q}(-k) \hat{q}(k) + \gamma_m^{(N)}(k) \hat{p}(-k) \hat{p}(k) \right) \\ &= \frac{1}{2} \left(\left\| (\gamma_m^{(N)})^{-1/2} \hat{q} \right\|_{\mathfrak{h}_{N,L}}^2 + \left\| (\gamma_m^{(N)})^{1/2} \hat{p} \right\|_{\mathfrak{h}_{N,L}}^2 \right), \end{aligned} \quad (4.5)$$

⁸We drop the dependence of the state $\omega_{L,0}^{(N)}$ on m , because we do not change $m > 0$ during the paper.

and hence $\omega_{\mu_N} = \omega_{L,0}^{(N)}$, and their GNS representations are the same. The expression (4.5) facilitates its relation to states on the continuum, see next Section.

The lattice “mass” μ_N is a dimensionless parameter and one can consider its “flow” with respect to the action of renormalization group (2.5) on states and Figure 2, cf. [82], in the following sense: We fix $\mu_N = \mu$ coherently along all scales and follow the variation of the parameter μ_{N+M} entering the renormalized states $\omega_{L,M}^{(N)}$ as a function of M , i.e. $\mu = \mu(M)$.

Although, strictly speaking, the “mass” parameter μ_{N+M} is not of the type as the initial “mass” μ_N because the scaling map α_{N+M}^N modifies the form of the dispersion relation and, thereby, the lattice interactions (if we were to interpret $\omega_{L,M}^{(N)}$ as ground state itself). This could be compensated by separating the part of $\omega_{L,M}^{(N)}$ that resembles the initial $\omega_{L,0}^{(N)}$, which would depend on μ_{N+M} , in addition to $\prod_{n=1}^M m_0(\varepsilon_{N+n})$ (see Section 4.2.2).

In order to obtain a convergent sequence of states as we scale ε_N , the dependence of μ_N on this UV cutoff must be fixed in such a way that it determines a finite physical mass m in the infrared.

4.1.2 Continuum free scalar field

Let us next fix the notations about the free scalar field on the continuum, both on the cylinder and on Minkowski spacetime (i.e., for the finite and infinite volume cases respectively). We pick a mass $m > 0$.

On the cylinder. We will regard the torus $\mathbb{T}_L^d = [-L, L]^d$ as the time zero Cauchy surface $\mathbb{T}_L^d \simeq \{0\} \times \mathbb{T}_L^d$ of the cylinder spacetime $\mathbb{R} \times \mathbb{T}_L^d$ endowed with the Minkowski metric. The Fourier transform of a function ξ on \mathbb{T}_L^d is defined as

$$\hat{\xi}(k) = \int_{[-L,L]^d} \xi(x) e^{-ixk} dx, \quad k \in \frac{\pi}{L} \mathbb{Z}^d. \quad (4.6)$$

The one-particle Hilbert space \mathfrak{h}_L of the mass m free scalar field on the cylinder is the completion of $C^\infty(\mathbb{T}_L^d, \mathbb{C})$ with the scalar product defined as follows: let $\xi, \eta \in C^\infty(\mathbb{T}_L^d, \mathbb{C})$ whose real and imaginary parts are denoted by $\xi = q_\xi + ip_\xi$ and $\eta = q_\eta + ip_\eta$, then their scalar product is

$$\langle \xi, \eta \rangle_L = \frac{1}{(2L)^d} \sum_{k \in \frac{\pi}{L} \mathbb{Z}^d} \overline{\left(\gamma_m^{-1/2}(k) \hat{q}_\xi(k) + i \gamma_m^{1/2}(k) \hat{p}_\xi(k) \right)} \left(\gamma_m^{-1/2}(k) \hat{q}_\eta(k) + i \gamma_m^{1/2}(k) \hat{p}_\eta(k) \right), \quad (4.7)$$

where $\gamma_m(k) = (k^2 + m^2)^{\frac{1}{2}}$ is the continuum dispersion relation. The complex structure (“multiplication by the imaginary unit”) on \mathfrak{h}_L is defined as

$$q + ip \longmapsto -\gamma_m p + i \gamma_m^{-1} q,$$

where $\widehat{(\gamma_m \xi)}(k) = \gamma_m(k) \hat{\xi}(k)$. The associated symplectic form is

$$\sigma_L(\xi, \eta) := \mathfrak{S}\langle \xi, \eta \rangle_L = \mathfrak{S} \left(\int_{[-L,L]^d} d^d x \bar{\xi}(x) \eta(x) \right). \quad (4.8)$$

The Fock representation π_L of the Weyl algebra $\mathcal{W}(\mathfrak{h}_L)$ is the GNS representation induced by the state

$$\omega_L(W_{\text{ct}}(\xi)) = e^{-\frac{1}{4}\|\xi\|_L^2}, \quad \xi \in \mathfrak{h}_L. \quad (4.9)$$

It will be convenient in the following not to consider π_L as acting on the Fock space built on \mathfrak{h}_L . Instead, we will realize it on the (mass independent) Fock space $\mathfrak{F}_+(L^2(\mathbb{T}_L^d)) \cong \mathfrak{F}_+(\ell^2(\frac{\pi}{L}\mathbb{Z}, (2L)^{-d}))$ by⁹

$$\pi_L(W_{\text{ct}}(\xi)) = e^{i\left[a\left(\gamma_m^{-1/2}q_\xi + i\gamma_m^{1/2}p_\xi\right) + a^*\left(\gamma_m^{-1/2}q_\xi + i\gamma_m^{1/2}p_\xi\right)\right]}, \quad \xi \in \mathfrak{h}_L, \quad (4.10)$$

where a and a^* are the standard creation and annihilation operator on $\mathfrak{F}_+(L^2(\mathbb{T}_L^d))$. Indeed, it is easy to check that the Fock vacuum $\Omega_L \in \mathfrak{F}_+(L^2(\mathbb{T}_L^d))$ is cyclic for the linear span of the operators (4.10) and induces the state ω_L .

Let S be an open subregion of \mathbb{T}_L^d , the (time-zero) local C^* -algebra associated to S is the C^* -subalgebra $\mathcal{W}_L(S)$ of $\mathcal{W}(\mathfrak{h}_L)$ generated by the Weyl operators $W_{\text{ct}}(\xi)$ with $\xi \in C^\infty(\mathbb{T}_L^d, \mathbb{C})$ compactly supported in S , and let $\mathcal{A}_L(S) := \pi_L(\mathcal{W}_L(S))''$ be the (time-zero) local von Neumann algebra associated to S in the representation.

The free field dynamics satisfies the Klein-Gordon equation. It is realized on \mathfrak{h}_L by a one parameter group of unitaries $\{\tau_{L|t}\}_{t \in \mathbb{R}}$:

$$\tau_{L|t}(\hat{\xi}) = [\cos(t\gamma_m) + i\gamma_m^{-1}\sin(t\gamma_m)]\hat{q}_\xi + i[\cos(t\gamma_m) + i\gamma_m\sin(t\gamma_m)]\hat{p}_\xi, \quad t \in \mathbb{R}. \quad (4.11)$$

The unitaries $\{\tau_{L|t}\}$, leaving the symplectic form invariant, induce a one parameter group $\{\eta_{L|t}\}$ of automorphisms of $\mathcal{W}(\mathfrak{h}_L)$ for which ω_L is an invariant state, and which are therefore implemented in the representation π_L by unitaries $U_L(0, t)$, $t \in \mathbb{R}$, on $\mathfrak{F}_+(L^2(\mathbb{T}_L^d))$. Moreover, as the Klein-Gordon equation has the speed of propagation 1, we have $\text{Ad } U_L(0, t)(\mathcal{A}_L(\mathcal{B}_r)) \subset \mathcal{A}_L(\mathcal{B}_{r+|t|})$, where \mathcal{B}_r is a ball of radius $r > 0$ and $r + |t| < L$. Similarly, the spacelike translations (on \mathfrak{h}_L) are unitarily implemented on $\mathfrak{F}_+(\mathfrak{h}_L)$ by $U_L(a, 0)$, $a \in \mathbb{R}^d$, and $U_L(a, 0), U_L(0, t)$ commute.

Altogether, $(\mathcal{A}_L, U_L, \Omega_L)$ form a continuum finite volume time-zero net of observables as per Section 2.4.

On Minkowski space. The construction is parallel to the cylinder case. Our convention for the Fourier transform of a function $\xi : \mathbb{R}^d \rightarrow \mathbb{C}$ is

$$\hat{\xi}(k) = \int_{\mathbb{R}^d} \xi(x) e^{-ixk} dx. \quad (4.12)$$

We note that with this convention and the one (4.6) for the torus, if ξ is a function compactly supported in $(-L, L)^d$ then the value of $\hat{\xi}(k)$ on $k \in \frac{\pi}{L}\mathbb{Z}^d$ is defined without ambiguity whether we consider ξ on the torus or on \mathbb{R}^d . This will be useful in the following.

The one-particle Hilbert space \mathfrak{h}_∞ for the continuum free field on Minkowski space is the completion of $C_0^\infty(\mathbb{R}^d, \mathbb{C})$ with respect to the scalar product defined for $\xi, \eta \in C_0^\infty(\mathbb{R}^d, \mathbb{C})$, (whose real and imaginary parts are $\xi = q_\xi + ip_\xi$ and $\eta = q_\eta + ip_\eta$) as

$$\langle \xi, \eta \rangle_\infty = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} dk \overline{\left(\gamma_m^{-1/2}(k)\hat{q}_\xi(k) + i\gamma_m^{1/2}(k)\hat{p}_\xi(k) \right)} \left(\gamma_m^{-1/2}(k)\hat{q}_\eta(k) + i\gamma_m^{1/2}(k)\hat{p}_\eta(k) \right), \quad (4.13)$$

⁹Here L^2 denotes the space of square-summable functions and not the square of the length L .

with the same dispersion relation as above. Also \mathfrak{h}_∞ becomes a complex Hilbert space if the complex structure is given by

$$q + ip \longmapsto -\gamma_m p + i\gamma_m^{-1} q,$$

and the associated symplectic form is

$$\sigma_\infty(\xi, \eta) := \Im \langle \xi, \eta \rangle_\infty = \Im \left(\int_{\mathbb{R}^d} d^d x \bar{\xi}(x) \eta(x) \right). \quad (4.14)$$

The Fock representation π_∞ of the Weyl algebra $\mathcal{W}(\mathfrak{h}_\infty)$ is the GNS representation specified by the state

$$\omega_\infty(W_{\text{ct}}(\xi)) = e^{-\frac{1}{4}\|\xi\|^2}, \quad \xi \in \mathfrak{h}_\infty, \quad (4.15)$$

and it is realized on the Fock space over $L^2(\mathbb{R}^d)$ (independent of m) by

$$\pi_\infty(W_{\text{ct}}(\xi)) = e^{i \left[a \left(\gamma_m^{-1/2} q_\xi + i\gamma_m^{1/2} p_\xi \right) + a^* \left(\gamma_m^{-1/2} q_\xi + i\gamma_m^{1/2} p_\xi \right) \right]}.$$

The (time-zero) local C^* -algebra $\mathcal{W}_\infty(S)$ and von Neumann algebra $\mathcal{A}_\infty(S)$ associated to bounded open regions $S \subset \mathbb{R}^d$ are defined similarly to the torus case considered above, and the dynamics is given by unitaries $\{\tau_{\infty|t}\}_{t \in \mathbb{R}}$ on \mathfrak{h}_∞ , whose action on a generic $\xi \in \mathfrak{h}_\infty$ is defined again by formula (4.11) and which again induce automorphisms of $\mathcal{W}(\mathfrak{h}_\infty)$ with finite propagation speed, implemented in π_∞ by unitaries $U_\infty(0, t)$, $t \in \mathbb{R}$. Thus, together with the unitary implementers $U_\infty(a, 0)$, $a \in \mathbb{R}^d$, of spatial translations, and with the Fock vacuum $\Omega_\infty \in \mathfrak{F}_+(L^2(\mathbb{R}^d))$, we obtain the continuum infinite volume time-zero net of observables $(\mathcal{A}_\infty, U_\infty, \Omega_\infty)$ defined by the free scalar field on Minkowski space.

We summarize in the following table our notations for one particle spaces, algebras, Weyl operators and vacuum states on the lattices and for the inductive limit and the continuum theories.

| | lattice object | inductive limit | continuum object (possibly $L = \infty$) |
|-------------------------|------------------------|---|--|
| one-particle space | $\mathfrak{h}_{N,L}$ | $\mathfrak{h}_{\infty,L}$ (symplectic space) | \mathfrak{h}_L |
| full algebra | $\mathcal{W}_{N,L}$ | $\mathcal{W}_{\infty,L}$ | \mathcal{W}_L |
| local algebra | $\mathcal{W}_{N,L}(S)$ | $\mathcal{W}_{\infty,L}(S)$ | $\mathcal{W}_L(S)$ |
| field | $\Phi_N(x)$ | | $\Phi(x)$ |
| momentum | $\Pi_N(x)$ | | $\Pi(x)$ |
| Weyl operator | $W_N(\xi)$ | | $W_{\text{ct}}(\xi)$ |
| state (of mass m) | $\omega_{L,M}^{(N)}$ | $\omega_{L,\infty}^{(\infty)}$ | ω_L |
| dispersion relation | $\gamma_m^{(N)}$ | | γ_m |

4.2 Constructing the continuum limit

In this section we will discuss the relation between the quasi-local and the local algebras of the inductive limit and of the continuum free field.

4.2.1 Embedding the lattice algebras into the continuum algebra

The lattice algebras in different scales with volume $(2L)^d$ are embedded into one another by $\alpha_{N+1}^N : \mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N+1,L}$. Here we show that they are further embedded into the continuum algebra $\mathcal{W}(\mathfrak{h}_L)$. This is realized by the \mathbb{R} -linear map $R_\infty^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_L$ defined by

$$R_\infty^N(\xi)(y) = \varepsilon_N^{\frac{d}{2}} \sum_{x \in \Lambda_N} (q(x) + ip(x)) \phi_x^{(\varepsilon_N)}(y), \quad (4.16)$$

where we recall that $q := \varepsilon_N^{-\frac{1+d}{2}} \Re \xi$, $p := \varepsilon_N^{\frac{1-d}{2}} \Im \xi$.

Proposition 4.1. *There exists an injective *-homomorphism $\beta_L : \mathcal{W}_{\infty,L} \rightarrow \mathcal{W}(\mathfrak{h}_L)$ such that*

$$\beta_L(W_N(\xi)) = W_{\text{ct}}(R_\infty^N(\xi)), \quad \xi \in \mathfrak{h}_{N,L}.$$

Proof. The map R_∞^N preserves the symplectic forms on $\mathfrak{h}_{N,L}$ and \mathfrak{h}_L , defined in Sections 3.1.2 and 4.1.2 respectively, indeed:

$$\begin{aligned} & \sigma_L(R_\infty^N(\xi), R_\infty^N(\eta)) \\ &= \Im \left(\varepsilon_N^d \sum_{x, x' \in \Lambda_N} \overline{(q_\xi(x) + ip_\xi(x)(q_\eta(x') + ip_\eta(x'))) \int_{[-L,L]^d} d^d y \varepsilon_N^{-d} \phi\left(\frac{y-x}{\varepsilon_N}\right) \phi\left(\frac{y-x'}{\varepsilon_N}\right)} \right) \\ &= \varepsilon_N^d \sum_{x \in \Lambda_N} q_\xi(x)p_\eta(x) - p_\xi(x)q_\eta(x) = \sigma_{N,L}(\xi, \eta), \end{aligned}$$

where the second equality holds since $\langle \phi_x^{(\varepsilon_N)}, \phi_{x'}^{(\varepsilon_N)} \rangle_L = \delta_{xx'}$ by the property of the scaling function ϕ . Therefore by the uniqueness of the Weyl C^* -algebra, R_∞^N induces an injective *-homomorphism of C^* -algebras $\beta_L^N : \mathcal{W}_{N,L} \rightarrow \mathcal{W}(\mathfrak{h}_L)$ such that $\beta_L^N(W_N(\xi)) = W_{\text{ct}}(R_\infty^N(\xi))$. Moreover, there holds

$$\begin{aligned} R_\infty^{N+1}(R_{N+1}^N(\xi)) &= \left(\frac{\varepsilon_N}{2}\right)^{\frac{d}{2}} \sum_{x \in \Lambda_{N+1}} (q_{R_{N+1}^N(\xi)}(x) + ip_{R_{N+1}^N(\xi)}(x)) \phi_x^{(\varepsilon_N/2)} \\ &= \left(\frac{\varepsilon_N}{2}\right)^{\frac{d}{2}} 2^{\frac{d}{2}} \sum_{y \in \Lambda_N} (q(y) + ip(y)) \sum_{n \in \mathbb{Z}^d} h_n \sum_{x \in \Lambda_{N+1}} \delta_{y+\frac{\varepsilon_N n}{2}}^{(\varepsilon_N/2)}(x) \phi_x^{(\varepsilon_N/2)} \\ &= \varepsilon_N^{\frac{d}{2}} \sum_{y \in \Lambda_N} (q(y) + ip(y)) \sum_{n \in \mathbb{Z}^d} h_n \phi_{y+\frac{\varepsilon_N n}{2}}^{(\varepsilon_N/2)} \\ &= \varepsilon_N^{\frac{d}{2}} \sum_{y \in \Lambda_N} (q(y) + ip(y)) \phi_y^{(\varepsilon_N)} = R_\infty^N(\xi), \end{aligned}$$

where in the fourth equality we used (3.39). This entails $\beta_L^{N+1} \circ \alpha_{N+1}^N = \beta_L^N$, and therefore, by the uniqueness and simplicity of the C^* -inductive limit, β_L exists as in the statement. \square

Now that we have realized the lattice algebras $\mathcal{W}_{N,L}$ inside $\mathcal{W}(\mathfrak{h}_L)$, we can consider the restrictions, $\omega_L|_{\beta_L(\mathcal{W}_{N,L})}$, of the vacuum state ω_L and take these as the “initial states”. This yields in a straightforward way a continuum time-zero net of local algebras. However, as already remarked, this choice is possible only because we know explicitly the vacuum state of the continuum free field. Below we study a more interesting case where we take the ground state (4.3) of the lattice free field at each scale, $\omega_{L,0}^{(N)} = \omega_{\mu_N}$, and show that the renormalization group scheme of Section 2.1 constructs ω_L as the scaling limit.

4.2.2 Scaling limits of free-field ground states

Following the general scheme of Section 2.1, the scaling limit of (4.3) is obtained by a two-step procedure, which we will explicitly implement:

1. Prove the convergence of $\lim_{M \rightarrow \infty} \omega_{L,M}^{(N)} = \omega_{L,\infty}^{(N)}$ on $\mathcal{W}_{N,L}$ at every fixed scale $N \in \mathbb{N}_0$.
2. Construct the projective-limit state $\varprojlim_N \omega_{L,\infty}^{(N)} = \omega_{L,\infty}^{(\infty)}$ on $\mathcal{W}_{\infty,L}$.

According to Figure 2, the sequence of states $\{\omega_{L,M}^{(N)}\}_{M \in \mathbb{N}_0}$ at each level N is generated by iterating the *flow equation* (2.5). In the case at hand, we use (3.29) and (3.10) to define the renormalization group, $\{\alpha_{N'}^N\}_{N < N'}$, resulting in:

$$\begin{aligned} & \omega_{L,M}^{(N)}(W_N(\xi)) \\ &= \omega_{\mu_{N+M},M}(W_N(\xi)) \\ &= e^{-\frac{2^{dM}}{4} \left(\left\| \gamma_{\mu_{N+M}}^{-1/2} \prod_{n=1}^M m_0(\varepsilon_{N+n} \cdot) \hat{q}_{N+M} \right\|_{\mathfrak{h}_{N+M}}^2 + \left\| \gamma_{\mu_{N+M}}^{1/2} \prod_{n=1}^M m_0(\varepsilon_{N+n} \cdot) \hat{p}_{N+M} \right\|_{\mathfrak{h}_{N+M}}^2 \right)}, \end{aligned} \quad (4.17)$$

for any $\xi \in \mathfrak{h}_{N,L}$, where we made the scale dependence of one-particle vectors explicit and denoted by $\hat{q}_{N+M}, \hat{p}_{N+M}$ the periodic extensions of \hat{q}_N, \hat{p}_N to Γ_{N+M} . Thus, if we choose $\{\mu_N\}$ which satisfies the *renormalization condition*,

$$\lim_{N \rightarrow \infty} \varepsilon_N^{-2} (\mu_N^2 - 2d) = m^2, \quad (4.18)$$

for a fixed ‘‘physical’’ mass $m > 0$, we find the free continuum dispersion relation of mass m in the scaling limit: $\lim_{M \rightarrow \infty} \gamma_{\mu_{N+M}}(k)^2 = m^2 + k^2 = \gamma_m(k)^2$. By (3.30), the limit state should be formally

$$\omega_{L,\infty}^{(N)}(W_N(\xi)) = e^{-\frac{1}{4} \left\| \hat{\phi}_0^{(\varepsilon_N)} (\gamma_m^{-1/2} \hat{q}_\infty + i \gamma_m^{1/2} \hat{p}_\infty) \right\|_{2,L}^2}, \quad (4.19)$$

where $\hat{q}_\infty, \hat{p}_\infty$ are the periodic extensions of \hat{q}_N, \hat{p}_N to the infinite lattice $\Gamma_\infty = \frac{\pi}{L} \mathbb{Z}^d$, and $\|\cdot\|_{2,L}$ is the standard norm on $\ell^2(\frac{\pi}{L} \mathbb{Z}^d, (2L)^{-d} \mu_{\Gamma_\infty})$. The norms involved there are finite if the scaling function decays like $|\hat{\phi}(k)| \leq C(1 + |k|)^{-\rho}$ with $\rho > \frac{d+1}{2}$. Our choice of the Daubechies wavelet family, $\{\phi_K\}_{K \in \mathbb{N}}$ [23] (see Section 3.3.4), is indeed sufficient if $\phi = {}_2\phi^{\otimes d}$ because $2\rho \approx 1.339$. Moreover, (4.19) is not only the formal limit of the sequence (4.17) but a weak*-limit point as the following lemma shows.

Lemma 4.2. *Assume that the renormalization condition (4.18) holds and that ϕ is built via tensor products from the Daubechies family, $\phi = {}_K\phi^{\otimes d}$ for $K \geq 2$. Then, the sequence of states $\{\omega_{L,M}^{(N)}\}_{M \in \mathbb{N}_0}$ on $\mathcal{W}_{N,L}$ is weak* convergent to the state (4.19) for every $N \in \mathbb{N}_0$.*

Proof. We state the proof for $\phi = {}_K\phi^{\otimes d}$ with $K \geq 2$.

Since the Weyl elements $W_N(\xi)$, $\xi \in \mathfrak{h}_{N,L}$, form a total set in $\mathcal{W}_{N,L}$ in norm, it is sufficient to prove the convergence of (4.17) to (4.19). To this end, we rewrite the $\|\cdot\|_{\mathfrak{h}_{N+M}}$ norms in (4.17) in terms of the norm $\|\cdot\|_{2,L}$ to be able to apply Lebesgue’s dominated convergence theorem for the measure space $(\frac{\pi}{L} \mathbb{Z}^d, (2L)^{-d} \mu_{\frac{\pi}{L} \mathbb{Z}^d})$:

$$\begin{aligned} & 2^{dM} \left\| \gamma_{\mu_{N+M}}^{1/2} \prod_{n=1}^M m_0(\varepsilon_{N+n} \cdot) \hat{p}_{N+M} \right\|_{\mathfrak{h}_{N+M}}^2 \\ &= \frac{\varepsilon_N^d}{(2L)^d} \sum_{k \in \frac{\pi}{L} \mathbb{Z}^d} \left| \chi_{\Gamma_{N+M}}(k) \gamma_{\mu_{N+M}}(k)^{1/2} \prod_{n=1}^M m_0(\varepsilon_{N+n} k) \hat{p}_{N+M}(k) \right|^2, \end{aligned}$$

where $\chi_S(k)$ is the characteristic function of the set S , and similarly for the term involving \hat{q}_{N+M} . Now, the periodic extensions $\hat{q}_{N+M}, \hat{p}_{N+M}$ of \hat{q}_N, \hat{p}_N are bounded in $k \in \Gamma_{N+M} \subset \Gamma_\infty = \frac{\pi}{L}\mathbb{Z}^d$, uniformly for $M \in \mathbb{N}$, and the same holds for $\gamma_{\mu_{N+M}}^{-1/2}$ thanks to the renormalization condition (4.18). As we know that

$$\left| \chi_{\Gamma_{N+M}}(k) \gamma_{\mu_{N+M}}(k)^\alpha \prod_{n=1}^M m_0(\varepsilon_{N+n} k) \right|^2, \quad -\frac{1}{2} \leq \alpha \leq \frac{1}{2} \quad (4.20)$$

converge pointwise to $|\hat{\phi}_0^{(\varepsilon_N)}(k) \gamma_m(k)^\alpha|^2$ by (3.30), we only have to show that this sequence is bounded by a $\mu_{\frac{\pi}{L}\mathbb{Z}^d}$ -integrable function uniformly for $M \in \mathbb{N}$. Next, we observe that because of the renormalization condition (4.18), we can replace

$$\gamma_{\mu_{N+M}}(k)^\alpha = \left(\varepsilon_{N+M}^{-2} (\mu_{N+M}^2 - 2d) + 2\varepsilon_{N+M}^{-2} \sum_{j=1}^d (1 - \cos(\varepsilon_{N+M} k_j)) \right)^{\frac{\alpha}{2}}$$

by

$$\left(m^2 + 2^{2M} \varepsilon_N^{-2} 2 \sum_{j=1}^d (1 - \cos(2^{-M} \varepsilon_N k_j)) \right)^{\frac{\alpha}{2}} = \left(m^2 + \sum_{j=1}^d (2^{M+1} \varepsilon_N^{-1} \sin(2^{-(M+1)} \varepsilon_N k_j))^2 \right)^{\frac{\alpha}{2}}.$$

Since we assume a tensor product structure for the scaling function, $\phi = {}_K\phi^{\otimes d}$, the same is true for the function m_0 , i.e. $m_0 = {}_K m_0^{\otimes d}$. Similarly, the characteristic function χ_{Γ_N} factorizes: $\chi_{\Gamma_N}(k) = \prod_{j=1}^d \chi_{\Gamma_N^{(1)}}(k_j)$ for $k \in \frac{\pi}{L}\mathbb{Z}^d$ with $\Gamma_N^{(1)}$ the $d = 1$ analogue of Γ_N . Therefore, we can use the estimate,

$$\begin{aligned} & \left(m^2 + \sum_{j=1}^d (2^{M+1} \varepsilon_N^{-1} \sin(2^{-(M+1)} \varepsilon_N k_j))^2 \right)^{\frac{\alpha}{2}} \\ & \leq \frac{\max(1, (\varepsilon_N m)^\alpha)}{\varepsilon_N^\alpha} \prod_{j=1}^d \left(1 + (2^{M+1} \sin(2^{-(M+1)} \varepsilon_N k_j))^2 \right)^{\frac{\alpha}{2}} \end{aligned}$$

to reduce the problem to the $d = 1$ case. Finally, to conclude the estimation of (4.20), we only need to find a $\mu_{\frac{\pi}{L}\mathbb{Z}}$ -integrable function that dominates the sequence:

$$\left| {}_K g_M^{(N)}(k) \right|^2 = \left| \chi_{\Gamma_{N+M}^{(1)}}(k) \left(1 + (2^{M+1} \sin(2^{-(M+1)} \varepsilon_N k))^2 \right)^{\frac{\alpha}{2}} \prod_{n=1}^M {}_K m_0(2^{-n} \varepsilon_N k) \right|^2.$$

In the following, we set $l = \varepsilon_N k \in \pi\mathbb{Z}$ to write:

$$\left| {}_K g_M^{(N)}(l) \right|^2 = \left| \chi_{\varepsilon_N \Gamma_{N+M}^{(1)}}(l) \left(1 + (2^{M+1} \sin(2^{-(M+1)} l))^2 \right)^{\frac{\alpha}{2}} \prod_{n=1}^M {}_K m_0(2^{-n} l) \right|^2. \quad (4.21)$$

Now, we observe that ${}_K m_0$ factorizes according to [23, (6.1.2)],

$${}_K m_0(l) = \left(\frac{1 + e^{-il}}{2} \right)^K {}_K \mathcal{L}(l),$$

where ${}_K\mathcal{L}$ is a certain trigonometric polynomial, which leads to:

$$\left| {}_K g_M^{(N)}(l) \right|^2 = \chi_{\varepsilon_N \Gamma_{N+M}^{(1)}}(l) \left(1 + (2^{M+1} \sin(2^{-(M+1)} l))^2 \right)^\alpha \prod_{n=1}^M |\cos(2^{-n} \frac{1}{2} l)|^{2K} |{}_K \mathcal{L}(2^{-n} l)|^2.$$

Using the analogue of Viète's (respectively Euler's) formula for finite products,

$$\prod_{n=1}^M \cos(2^{-n} \frac{1}{2} l) = \frac{2 \sin(\frac{1}{2} l)}{2^{M+1} \sin(2^{-(M+1)} l)},$$

(which is a consequence of iterating the cosine product identity, $\cos(\varphi) \cos(\vartheta) = \frac{1}{2}(\cos(\varphi + \vartheta) + \cos(\varphi - \vartheta))$, followed by a finite geometric summation¹⁰), we find:

$$\begin{aligned} & \left| {}_K g_M^{(N)}(l) \right|^2 \\ &= \chi_{\varepsilon_N \Gamma_{N+M}^{(1)}}(l) \left| \frac{2 \sin(\frac{1}{2} l)}{2^{M+1} \sin(2^{-(M+1)} l)} \right|^{2K} \left(1 + (2^{M+1} \sin(2^{-(M+1)} l))^2 \right)^\alpha \prod_{n=1}^M |{}_K \mathcal{L}(2^{-n} l)|^2, \end{aligned}$$

where in the second equality we used that if $l \in \varepsilon_N \Gamma_{N+M}^{(1)}$ then $2^{-(M+1)} |l| \leq \pi/2$. Next, we apply the basic inequalities,

$$\begin{aligned} 2^{M+1} \sin(2^{-(M+1)} |l|) &\leq |l|, & \forall l \in \mathbb{R}, \\ \frac{2}{\pi} |l| &\leq 2^{M+1} \sin(2^{-(M+1)} |l|), & \forall l \in \mathbb{R} : |l| \leq 2^M \pi, \end{aligned}$$

to arrive at:

$$\begin{aligned} \left| {}_K g_M^{(N)}(l) \right|^2 &\leq \chi_{\varepsilon_N \Gamma_{N+M}^{(1)}}(l) \pi^{2K} \left(\frac{|\sin(\frac{1}{2} l)|}{|l|} \right)^{2K} (1 + |l|^2)^\alpha \prod_{n=1}^M |{}_K \mathcal{L}(2^{-n} l)|^2 \\ &\leq \chi_{\varepsilon_N \Gamma_{N+M}^{(1)}}(l) \pi^{2K} \left(\frac{|\sin(\frac{1}{2} l)|}{|l|} \right)^{2K} (1 + |l|)^{2\alpha} \prod_{n=1}^M |{}_K \mathcal{L}(2^{-n} l)|^2. \end{aligned}$$

Thus, to dominate the sequence (4.21) by $\mu_{\mathbb{Z}}$ -integrable function, it is sufficient to have an estimate of the form,

$$\prod_{n=1}^M |{}_K \mathcal{L}(2^{-n} l)| \leq C(1 + |l|)^{K-1-\epsilon}, \quad (4.22)$$

for some $C > 0$ and arbitrary $\epsilon > 0$, as this implies:

$$\left| {}_K g_M^{(N)}(l) \right|^2 \leq C^2 \pi^{2K} \left(\frac{|\sin(\frac{1}{2} l)|}{|l|} \right)^{2K} (1 + |l|)^{2(K+(\alpha-\epsilon)-1)}.$$

But the estimate (4.22) is an immediate consequence of the regularity analysis for compactly supported wavelets in [23, Section 7.1]. To see this, we observe that according to [23, Lemma 7.1.4]:

$$\sup_l |{}_K \mathcal{L}(l)| < 2^{K-1}. \quad (4.23)$$

Now, applying the reasoning of [23, Lemma 7.1.1] to the finite product $\prod_{n=1}^M |{}_K \mathcal{L}(2^{-n} l)|$ for the momentum space shells $|l| \leq 1$ and $2^{n-1} < |l| \leq 2^n$, $n \in \mathbb{N}_{\leq M}$ as well as $2^M < |l| \leq 2^M \pi$, we deduce that (4.23) implies (4.22) for $\epsilon > 0$ sufficiently small. \square

¹⁰See https://en.wikipedia.org/wiki/Vi%C3%A8te%27s_formula.

Actually, with the embedding $\beta_L : \mathcal{W}_{\infty,L} \rightarrow \mathcal{W}(\mathfrak{h}_L)$, the expression (4.19) is exactly the evaluation of the usual continuum vacuum state ω_L of mass m (4.9) on the continuum Weyl operators $W_{\text{ct}}(R_{\infty}^N(\xi))$.

Lemma 4.3. *It holds that $\omega_{L,\infty}^{(N)}(W) = \omega_L(\beta_L(W))$ for $W \in \mathcal{W}_{N,L}$.*

Proof. The one-particle symplectic map R_{∞}^N , which satisfies $\beta_L(W_N(\xi)) = W_{\text{ct}}(R_{\infty}^N(\xi))$, intertwines the states: $\omega_L(W_{\text{ct}}(R_{\infty}^N(\xi))) = \omega_{L,\infty}^{(N)}(W_N(\xi))$, $\xi \in \mathfrak{h}_{N,L}$. Indeed, using (4.7), (4.9) and (4.19),

$$\begin{aligned}
\omega_L(W_{\text{ct}}(R_{\infty}^N(\xi))) &= \\
&= \exp \left\{ -\frac{1}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \left| \gamma_m^{-\frac{1}{2}}(k) (\Re R_{\infty}^N(\xi))^{\wedge}(k) + i\gamma_m^{\frac{1}{2}}(k) (\Im R_{\infty}^N(\xi))^{\wedge}(k) \right|^2 \right\} \\
&= \exp \left\{ -\frac{1}{2(2r\varepsilon_N)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \left(\frac{1}{\gamma_m(k)} \left| \varepsilon_N^{d/2} \sum_{x \in \Lambda_N} q(x) \int_{[-L,L]^d} d^d y \phi_x^{(\varepsilon_N)}(y) e^{-iky} \right|^2 \right. \right. \\
&\quad \left. \left. + \gamma_m(k) \left| \varepsilon_N^{d/2} \sum_{x \in \Lambda_N} p(x) \int_{[-L,L]^d} d^d y \phi_x^{(\varepsilon_N)}(y) e^{-iky} \right|^2 \right) \right\} \\
&= \exp \left\{ -\frac{1}{2(2r\varepsilon_N)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \left(\frac{\varepsilon_N^{2d}}{\gamma_m(k)} \left| \sum_{x \in \Lambda_N} q(x) e^{-ikx} \right|^2 \right. \right. \\
&\quad \left. \left. + \varepsilon_N^{2d} \gamma_m(k) \left| \sum_{x \in \Lambda_N} p(x) e^{-ikx} \right|^2 \right) |\hat{\phi}(\varepsilon_N k)|^2 \right\} \\
&= \omega_{L,\infty}^{(N)}(W_N(\xi))
\end{aligned}$$

where in the third equality the relation

$$\int_{[-L,L]^d} d^d y \phi_x^{(\varepsilon_N)}(y) e^{-iky} = \varepsilon_N^{\frac{d}{2}} e^{-ikx} \hat{\phi}(\varepsilon_N k),$$

for $k \in \frac{\pi}{L}\mathbb{Z}$ was used, where $\hat{\phi}(\varepsilon_N k) = \int_{[-L,L]^d} d^d y \phi(y) e^{-i\varepsilon_N k y}$. \square

The projective consistency (2.9) of the limit states $\omega_{L,\infty}^{(N)}$ follows both abstractly from Proposition 2.2 and directly by applying (3.31). Therefore, we conclude that the limit state $\omega_{L,\infty}^{(\infty)}$ exists on $\mathcal{W}_{\infty,L}$. By the preceding Lemma, $\omega_{L,\infty}^{(\infty)}$ agrees with the usual continuum vacuum state ω_L on the span of Weyl operators indexed by functions $f \in L^2(\mathbb{T}_L^d)$ with finite wavelet expansion (3.36) associated with ϕ .

Remark 4.4. *The construction of suitable scaling functions ϕ in dimension $d \geq 2$ by tensor products is probably not optimal: what is needed is the condition $|\hat{\phi}(k)| \leq C(1 + |k|)^{-\rho}$ with $\rho > \frac{d+1}{2}$ which is rotationally invariant, while our wavelets are regular in d directions separately.*

To summarize:

Proposition 4.5. *Let $\{\mu_N\}_N$ satisfy (4.18) with $m > 0$ and $\{\alpha_{N'}^N\}$ be the scaling map defined through the Daubechies scaling function $K\phi$ with $K \geq 2$ on the scalar lattice algebras $\{\mathcal{W}_{N,L}\}$ as in Section 3.3. Then the sequences of states $\{\omega_{L,M}^{(N)}\}_{M \in \mathbb{N}_0} = \{\omega_{\mu_{N+M},M}\}_{M \in \mathbb{N}_0}$ has the scaling limit $\omega_{L,\infty}^{(\infty)}$ on $\mathcal{W}_{\infty,L}$ in the sense of (2.11). With the embedding β_L of $\mathcal{W}_{\infty,L}$ in $\mathcal{W}(\mathfrak{h}_L)$ by Proposition 4.1, it holds that $\omega_{L,\infty}^{(\infty)}(W_N(\xi)) = \omega_L(\beta_L(W_N(\xi)))$ for $\xi \in \mathfrak{h}_{N,L}$ with some N .*

4.2.3 Density of the wavelet observables in the continuum limit

Let $\pi_{L,\infty}^{(\infty)}$ be the GNS representation of $\mathcal{W}_{\infty,L}$ w.r.t. $\omega_{L,\infty}^{(\infty)}$. Here we show that the local C^* -algebra $\mathcal{W}_{\infty,L}(S)$, defined as in (2.14), is dense in $\mathcal{W}_L(S)$ in the strong operator topology in the representation $\pi_{L,\infty}^{(\infty)}$, if we use a sufficiently regular scaling function (with $K \geq 6$). Recall that $\phi_x^{(\varepsilon_N)} \in L^2(\mathbb{T}_L^d)$, $x \in \Lambda_N$, is the $2L$ -periodization of the function $y \in \mathbb{R}^d \mapsto \varepsilon_N^{-\frac{d}{2}} \phi(\frac{y-x}{\varepsilon_N})$ restricted to an open cube of side length $2L$ containing its support.

Lemma 4.6. *Let $K \geq 6$. Then the (complex) linear span of the functions $\phi_x^{(\varepsilon_N)}$, $x \in \Lambda_N$, $N \in \mathbb{N}$, is dense in \mathfrak{h}_L .*

Proof. As the norms are equivalent, \mathfrak{h}_L can be identified, as a real Hilbert space, with the direct sum of real Sobolev spaces $H_{\mathbb{R}}^{-\frac{1}{2}}(\mathbb{T}_L^d) \oplus H_{\mathbb{R}}^{\frac{1}{2}}(\mathbb{T}_L^d)$. With the notation of Section 3.3, we then recall that if $f \in H_{\mathbb{R}}^{\pm\frac{1}{2}}(\mathbb{R}^d)$ then for any $\ell_0 \geq 0$,

$$f = \sum_{n \in \mathbb{Z}^d} \langle \phi_{\ell_0,n}, f \rangle \phi_{\ell_0,n} + \sum_{\ell \geq \ell_0} \sum_{n \in \mathbb{Z}^d} \langle \psi_{\ell,n}, f \rangle \psi_{\ell,n} \quad (4.24)$$

where $\langle \cdot, \cdot \rangle$ denotes the standard $L^2(\mathbb{R}^d)$ inner product, and this series converges also in $H_{\mathbb{R}}^{\pm\frac{1}{2}}$ for sufficiently regular wavelets. Convergence in $H_{\mathbb{R}}^{-\frac{1}{2}}(\mathbb{R}^d)$ comes from convergence in $L^2(\mathbb{R}^d)$ and convergence in $H_{\mathbb{R}}^{\frac{1}{2}}(\mathbb{R}^d)$ comes with $K \geq 6$ by [43, Corollary 9.1, Example 9.1] since the $B_{22}^{\frac{1}{2}}$ Besov norm is equivalent to the $H_{\mathbb{R}}^{\frac{1}{2}}(\mathbb{R}^d)$ norm.

Let $f \in H_{\mathbb{R}}^{\pm\frac{1}{2}}(\mathbb{R}^d)$ have support contained in $(-L+\delta, L-\delta)^d$ for some $\delta > 0$. Then, since $\text{supp } \phi_{\ell,n} = 2^{-\ell}(\text{supp } \phi + n)$, and analogously for $\psi_{\ell,n}$, we can find a sufficiently large ℓ_0 such that the non-zero components in the expansion (4.24) of the rescaled function $f_\varepsilon = f(\varepsilon \cdot)$ all have supports in $(-\varepsilon^{-1}(L-\delta/2), \varepsilon^{-1}(L-\delta/2))^d$. Moreover, there holds the further expansion (actually a finite linear combination)

$$\psi_{\ell,n} = \sum_{k \in \mathbb{Z}^d} \langle \phi_{\ell+1,k}, \psi_{\ell,n} \rangle \phi_{\ell+1,k},$$

and again due to the support properties of $\psi_{\ell,n}$ the only non-zero coefficients involve functions $\phi_{\ell+1,m}$ whose support is contained in $(-\varepsilon^{-1}L, \varepsilon^{-1}L)^d$. Summing up, we obtain that f_ε can be approximated, in the $H_{\mathbb{R}}^{\pm\frac{1}{2}}(\mathbb{R}^d)$ norm, by finite linear combinations of functions $\phi_{j,n}$ with support contained in $(-\varepsilon^{-1}L, \varepsilon^{-1}L)^d$. Recalling now that $\phi_x^{(\varepsilon_N)}$ is the $2L$ -periodization of $\varepsilon^{-d/2} \phi_{N, \varepsilon_N^{-1}x}(\varepsilon^{-1} \cdot)$, we conclude that f can be approximated in $H_{\mathbb{R}}^{\pm 1/2}(\mathbb{T}_L^d)$ by finite real linear combinations of the functions $\phi_x^{(\varepsilon_N)}$, $x \in \Lambda_N$, $N \geq \ell_0$.

Note that if $f \in H_{\mathbb{R}}^{\pm\frac{1}{2}}(\mathbb{T}_L^d)$ has a support contained in a shifted torus, then f can be approximated as above because the wavelet basis generated by $\{\phi_x^{(\varepsilon_N)} : x \in \Lambda_N, N \geq \ell_0\}$

(ℓ_0 depends on the length of the support of f) is invariant under dyadic translations. For a general $f \in H_{\mathbb{R}}^{\pm\frac{1}{2}}(\mathbb{T}_L^d)$, we can take a smooth partition of unity on \mathbb{T}_L^d such that each support is contained in a shifted torus. Then f can be written as a finite sum of functions whose supports are contained in a shifted torus, and each of them can be approximated in the H^s -norm, hence so can f itself. \square

At this point we can show that our wavelet scaling limit gives the time-zero algebras of the usual massive free field on the cylinder.

Theorem 4.7. *Let $K \geq 6$. Let $\pi_{L,\infty}^{(\infty)}$, π_L be the GNS representations of $\mathcal{W}_{\infty,L}$, $\mathcal{W}(\mathfrak{h}_L)$ with respect to $\omega_{L,\infty}^{(\infty)}$, ω_L , and β_L be the embedding $\mathcal{W}_{\infty,L} \rightarrow \mathcal{W}(\mathfrak{h}_L)$ as in Proposition 4.1. Then there is a unitary operator $V : \mathcal{H}_{L,\infty}^{(\infty)} \rightarrow \mathcal{H}_L$ such that*

$$V\pi_{L,\infty}^{(\infty)}(W_N(\xi))V^* = \pi_L(\beta_L(W_N(\xi))), \quad \xi \in \mathfrak{h}_{N,L}, N \in \mathbb{N}.$$

Furthermore, we have $V\pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L}(S))''V^* = \pi_L(\mathcal{W}_L(S))''$ for any open set $S \subset \mathbb{T}_L^d$, and $V^*U_L(a,0)V$, $a \in \bigcup_{N \in \mathbb{N}} \Lambda_N$, implements the spatial dyadic translations $\eta_{L|a}^{(\infty)}$ of $\mathcal{W}_{\infty,L}$ in the representation $\pi_{L,\infty}^{(\infty)}$, where $U_L(a,0)$, $a \in \mathbb{T}_L^d$, are the spatial translations for π_L .

Proof. By Lemma 4.6, and by the strong continuity of $\xi \mapsto \pi_L(W_{\text{ct}}(\xi))$, we obtain $\pi_L(\beta_L(\mathcal{W}_{\infty,L}))'' = \pi_L(\mathcal{W}(\mathfrak{h}_L))'' = \mathcal{B}(\mathcal{H}_L)$. The unitary V is given by the uniqueness of GNS representation induced by $\omega_{L,\infty}^{(\infty)}$, together with Lemma 4.3 and the cyclicity of the GNS vector for $\pi_L(\beta_L(\mathcal{W}_{\infty,L}))'' = \mathcal{B}(\mathcal{H}_L)$.

As for local density, it is immediate to see that for any open set $S \subset \mathbb{T}_L^d$,

$$V\pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L}(S))''V^* \subset \pi_L(\mathcal{W}_L(S))''.$$

Let $S \subset (-L, L)^d$ and consider $\xi \in \mathfrak{h}_L$ such that $\text{supp } \xi \subset S$. By arguing as in Lemma 4.6, we can choose N_0 large enough such that we can approximate ξ by finite linear combinations of scaling functions $\{\phi_x^{(\varepsilon N)} : \text{supp } \phi_x^{(\varepsilon N)} \subset S, N \geq N_0\}$. Moreover, as $R_\infty^N \delta_x^{(N)} = \varepsilon_N^{-\frac{1}{2}} \phi_x^{(\varepsilon N)}$, $\text{supp } \phi_x^{(\varepsilon N)} \subset S$ entails $W_N(\delta_x^{(N)}) \in \mathcal{W}_{\infty,L}(S)$ and therefore, by strong continuity of Weyl operators,

$$\pi_L(W_{\text{ct}}(\xi)) \in V\pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L}(S))''V^*,$$

hence we have the equality $V\pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L}(S))''V^* = \pi_L(\mathcal{W}_L(S))''$.

Let now $S \subsetneq \mathbb{T}_L^d$ be a general open region and $\xi \in \mathfrak{h}_L$ such that $\text{supp } \xi \subset S$. Again by the argument of Lemma 4.6, we can write ξ as a finite sum of functions in \mathfrak{h}_L each of which has support in an open subset of S which can be brought inside the fundamental domain $(-L, L)^d$ by a translation in $\varepsilon_{N_0} \mathbb{Z}^d$ with $N_0 \in \mathbb{N}$ sufficiently large. The proof is then concluded by the above argument, together with the fact that the set $\{\phi_x^{(\varepsilon N)} : x \in \Lambda_N, N \geq N_0\}$ is globally invariant under $\varepsilon_{N_0} \mathbb{Z}^d$ -translations.

Finally, it is clear that the map $R_\infty^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_L$ in Eq. (4.16) intertwines the action of the dyadic translations on the lattice and on the torus \mathbb{T}_L^d . Since β_L is induced by R_∞^N , this immediately entails the statement about the implementation of the dyadic translations by $V^*U_L(a,0)V$, $a \in \bigcup_{N \in \mathbb{N}} \Lambda_N$. \square

We stress the fact that by the above result, the dyadic translations of the inductive limit theory extend continuously to the whole torus translation group.

4.2.4 Limit of dynamics

The free lattice dynamics $\eta_L^{(N)} : \mathbb{R} \curvearrowright \mathcal{W}_{N,L}$ induced by the Hamiltonian $H_{L,0}^{(N)}$ (4.1) results via second quantization from the harmonic time evolution, $\tau_L^{(N)} : \mathbb{R} \curvearrowright \mathfrak{h}_{N,L}$, on the one-particle space (using the conventions of Section 3.1.2, see e.g. [4, (1.8.30)]). In the momentum-space representation, i.e. on $\hat{\mathfrak{h}}_{N,L}$, it is given by the multiplication operator:

$$\tau_{L|t}^{(N)}(\hat{\xi}) = [\cos(t\gamma_{\mu_N}) + i\varepsilon_N^{-1}\gamma_{\mu_N}^{-1}\sin(t\gamma_{\mu_N})]\Re\hat{\xi} + i[\cos(t\gamma_{\mu_N}) + i\varepsilon_N\gamma_{\mu_N}\sin(t\gamma_{\mu_N})]\Im\hat{\xi}, \quad t \in \mathbb{R}, \quad (4.25)$$

which is the analogue of the free (continuum) time evolution, $\tau_L : \mathbb{R} \curvearrowright \mathfrak{h}_L$, given in (4.11) using γ_{μ_N} instead of γ_m . As this evolution preserves the symplectic structure, $\sigma_{N,L} \circ \tau_{L|t}^{(N)} = \sigma_{N,L}$, we obtain $\eta_L^{(N)}$ via

$$\eta_{L|t}^{(N)}(W_N(\hat{q}, \hat{p})) = W_N(\tau_{L|t}^{(N)}(\hat{q}, \hat{p})), \quad t \in \mathbb{R}, \quad (4.26)$$

where we write $\xi \in \mathfrak{h}_{N,L}$ as the tuple (\hat{q}, \hat{p}) as before. By construction, each initial state is preserved by the dynamics of the same scale, i.e. $\omega_{\mu_N,0} \circ \eta_{L|t}^{(N)} = \omega_{\mu_N,0}$, but the dynamics induced by $H_{L,0}^{(N+1)}$ does not preserve the image of α_{N+1}^N . In other words, $\eta_L^{(N')}$ does not preserve $\alpha_{N'}^N(\mathcal{W}_{N,L}) \subset \mathcal{W}_{N',L}$, $N' > N$. In terms of the one-particle spaces this follows from the fact that multiplying a periodically extended tuple of vectors $(\hat{q}, \hat{p}) \in \ell^2(\Gamma_N, (2r_N)^{-d}\mu_{\Gamma_N})$ by a function only periodic on Γ_{N+1} does not preserve¹¹ the periodicity on Γ_N . Moreover, we need to analyze the convergence of the dynamics relative to inductive limit structure of $\mathcal{W}_{\infty,L}$ and the GNS representation of $\omega_{L,\infty}^{(\infty)}$ because the lattice dynamics $\eta_L^{(N)}$ does not extend¹² to the $\mathcal{W}(\mathfrak{h}_L)$ and the continuum dynamics η_L is only defined with respect to $\pi_{L,\infty}^{(\infty)}$ (since τ_L does not preserve the subspace $\mathfrak{h}_{\infty,L} \subset \mathfrak{h}_L$). To this end, we use (4.16) to realize the map α_{∞}^N explicitly. In the Fourier space, it is induced by

$$R_{\infty}^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_{\infty,L} \subset \mathfrak{h}_L, \quad R_{\infty}^N(\hat{q}, \hat{p}) = \varepsilon_N^{\frac{d}{2}}\hat{\phi}(\varepsilon_N \cdot)(\hat{q}, \hat{p}) \quad (4.27)$$

where the abstract inductive limit $\mathfrak{h}_{\infty,L} = \varinjlim_N \mathfrak{h}_{N,L}$ was identified with the subspace of \mathfrak{h}_L spanned by functions in the range of the maps R_{∞}^N , $N \in \mathbb{N}$ thanks to the consistency $R_{\infty}^{N'} \circ R_{N'}^N = R_{\infty}^N$ for $N' > N$ (shown in the proof of Proposition 4.1) and the uniqueness of the inductive limit. This allows us to investigate the convergence of the sequence of dynamics according to the general form given in (2.13) which can be facilitated at the level of one-particle spaces:

Proposition 4.8. *For $N \in \mathbb{N}_0$ and for $(\hat{q}, \hat{p}) \in \mathfrak{h}_{N,L}$, we have:*

$$\left\| R_{\infty}^{N'}(\tau_{L|t}^{(N')}(R_{N'}^N(\hat{q}, \hat{p}))) - \tau_{L|t}(R_{\infty}^N(\hat{q}, \hat{p})) \right\|_{\mathfrak{h}_L} \rightarrow 0, \quad \text{as } N' \rightarrow \infty,$$

uniformly on compact sets of $t \in \mathbb{R}$.

¹¹See also Section 5.3 for a scaling map scheme using sharp momentum cutoffs, where such a consistent dynamics induced by $H_{L,0}^{(N+1)}$ relative to α_{N+1}^N exists.

¹²In terms of the one-particle spaces, this issue manifests in the following way: The complete space \mathfrak{h}_L is isomorphic as real topological vector space to $H^{-\frac{1}{2}}(\mathbb{T}_L^d) \oplus H^{\frac{1}{2}}(\mathbb{T}_L^d)$ which makes it evident that the dynamics $\tau_L^{(N)}$ does not extend to \mathfrak{h}_L because $\mathcal{F}^{-1}[\gamma_{\mu_N}^{-1}\hat{g}] \notin H^{\frac{1}{2}}(\mathbb{T}_L^d)$ for $g \in H^{-\frac{1}{2}}(\mathbb{T}_L^d)$.

Proof. As $\gamma_{\mu_{N'}} \rightarrow \gamma_m$ pointwise by (4.18), we use dominated convergence relative to the measure space $(\frac{\pi}{L}\mathbb{Z}^d, (2L)^{-d} \mu_{\frac{\pi}{L}\mathbb{Z}^d})$ and the decay properties of $\hat{\phi}$. Moreover, by (3.29)(3.30) we have an integrability estimate uniform in $t \in \mathbb{R}$:

$$\begin{aligned}
& \left\| R_\infty^{N'}(\tau_{L|t}^{(N')}(R_{N'}^N(\hat{q}, \hat{p}))) - \tau_{L|t}(R_\infty^N(\hat{q}, \hat{p})) \right\|_{\mathfrak{h}_L}^2 \\
&= \frac{\varepsilon_N^d}{(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} |\hat{\phi}(\varepsilon_N k)|^2 \left(\gamma_m(k)^{-1} |(\cos(\gamma_{\mu_{N'}}(k)t) - \cos(\gamma_m(k)t))\hat{q}(k) \right. \\
&\quad \left. - (\gamma_{\mu_{N'}}(k) \sin(\gamma_{\mu_{N'}}(k)t) - \gamma_m(k) \sin(\gamma_m(k)t))\hat{p}(k) \right|^2 \\
&\quad + \gamma_m(k) |(\cos(\gamma_{\mu_{N'}}(k)t) - \cos(\gamma_m(k)t))\hat{p}(k) \\
&\quad \left. + (\gamma_{\mu_{N'}}(k)^{-1} \sin(\gamma_{\mu_{N'}}(k)t) - \gamma_m(k)^{-1} \sin(\gamma_m(k)t))\hat{q}(k) \right|^2 \\
&\leq \frac{4\varepsilon_N^d}{(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \gamma_m(k) |\hat{\phi}(\varepsilon_N k)|^2 \left((\gamma_m(k)^{-1} |\hat{q}(k)| + (1+c')^{\frac{1}{2}} |\hat{p}(k)|)^2 \right. \\
&\quad \left. + \left(|\hat{p}(k)| + \frac{(1-c'')^{-\frac{1}{2}} + 1}{2} m^{-1} |\hat{q}(k)| \right)^2 \right) \\
&< \infty,
\end{aligned}$$

where from the second to the third line, we use the fact that the integrand in the second line is pointwise dominated by the integrand in the third line. This estimate uses the basic inequalities $(1-c'')^{\frac{1}{2}} m \leq \gamma_{\mu_{N'}}$ and $\gamma_{\mu_{N'}} \leq (1+c')^{\frac{1}{2}} \gamma_m$ valid uniformly in $k \in \frac{\pi}{L}\mathbb{Z}^d$ for sufficiently large $N' \in \mathbb{N}_0$ and constants $0 < c', c'' < 1$. Now, we can exploit the uniform continuity of \cos and \sin to conclude that the convergence is uniform on compact sets of $t \in \mathbb{R}$. More precisely, we have for $|t| \leq T < \infty$:

$$\begin{aligned}
& \sup_{|t| \leq T} \left\| R_\infty^{N'}(\tau_{L|t}^{(N')}(R_{N'}^N(\hat{q}, \hat{p}))) - \tau_{L|t}(R_\infty^N(\hat{q}, \hat{p})) \right\|_{\mathfrak{h}_L}^2 \\
&\leq \frac{\varepsilon_N^d}{(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} |\hat{\phi}(\varepsilon_N k)|^2 \left(\sup_{|t| \leq T} \gamma_m(k)^{-1} |(\cos(\gamma_{\mu_{N'}}(k)t) - \cos(\gamma_m(k)t))\hat{q}(k) \right. \\
&\quad \left. - (\gamma_{\mu_{N'}}(k) \sin(\gamma_{\mu_{N'}}(k)t) - \gamma_m(k) \sin(\gamma_m(k)t))\hat{p}(k) \right|^2 \\
&\quad + \sup_{|t| \leq T} \gamma_m(k) |(\cos(\gamma_{\mu_{N'}}(k)t) - \cos(\gamma_m(k)t))\hat{p}(k) \\
&\quad \left. + (\gamma_{\mu_{N'}}(k)^{-1} \sin(\gamma_{\mu_{N'}}(k)t) - \gamma_m(k)^{-1} \sin(\gamma_m(k)t))\hat{q}(k) \right|^2
\end{aligned}$$

Now, because the preceding estimate is independent of t , we can use the dominated convergence theorem, the fact that the pointwise supremum of a sequence of measurable function is again measurable, and that for every $k \in \frac{\pi}{L}\mathbb{Z}^d$,

$$\begin{aligned}
& \lim_{N' \rightarrow \infty} \sup_{|t| \leq T} |\cos(\gamma_{\mu_{N'}}(k)t) - \cos(\gamma_m(k)t)| = 0, \\
& \lim_{N' \rightarrow \infty} \sup_{|t| \leq T} |\gamma_{\mu_{N'}}(k)^{-1} \sin(\gamma_{\mu_{N'}}(k)t) - \gamma_m(k)^{-1} \sin(\gamma_m(k)t)| = 0
\end{aligned}$$

by the uniform continuity of \sin and \cos , to conclude the proof. \square

With $V : \mathcal{H}_{L,\infty}^{(\infty)} \rightarrow \mathcal{H}_L$ the unitary operator in Thm. 4.7 and $\{\eta_{L|t}\}$ the automorphisms of $\mathcal{W}(\mathfrak{h}_L)$ realizing the free dynamics, it is clear that

$$U_L(0, t) V \pi_{L,\infty}^{(\infty)}(W) V^* U_L(0, t) = \pi_L(\eta_{L|t}(\beta_L(W))), \quad W \in \mathcal{W}_{\infty, L},$$

so we can use V to transport the continuum free dynamics on $\mathcal{H}_{L,\infty}^{(\infty)}$ as $U_{L|t} := V^*U_L(0,t)V$, and by a slight abuse of notation we will also use $\eta_{L|t}$ to denote $\text{Ad}_{U_{L|t}}$.

Corollary 4.9. *Given $N \in \mathbb{N}_0$, $\{\pi_{L,\infty}^{(\infty)}(\alpha_{\infty}^{N'}(\eta_{L|t}^{(N')}(\alpha_{N'}^N(W))))\}_{N'>N}$ converges strongly to $\eta_{L|t}(\pi_{L,\infty}^{(\infty)}(\alpha_{\infty}^N(W)))$ for all $W \in \mathcal{W}_{N,L}$ and uniformly on compact sets of $t \in \mathbb{R}$.*

Proof. This will follow from Proposition 4.8 if we prove that Weyl operators in the representation $\pi_{L,\infty}^{(\infty)}$ viewed as maps from the Hilbert space \mathfrak{h}_L to $B(\mathcal{H}_L)$ are Lipschitz maps, where $B(\mathcal{H}_L)$ is given the strong operator topology. To this end, we first observe that by Theorem 4.7,

$$\begin{aligned} & \left\| \left[\pi_{L,\infty}^{(\infty)}(\alpha_{\infty}^{N'}(\eta_{L|t}^{(N')}(\alpha_{N'}^N(W)))) - \eta_{L|t}(\pi_{L,\infty}^{(\infty)}(\alpha_{\infty}^N(W))) \right] \pi_{L,\infty}^{(\infty)}(W_M(\zeta))\Omega_{L,\infty}^{(\infty)} \right\| \\ &= \left\| \left[\pi_L(\beta_L(\alpha_{\infty}^{N'}(\eta_{L|t}^{(N')}(\alpha_{N'}^N(W)))) - \pi_L(\eta_{L|t}(\beta_L(\alpha_{\infty}^N(W)))) \right] \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L \right\| \end{aligned}$$

for $\zeta \in \mathfrak{h}_{M,L}$ and some $M \in \mathbb{N}_0$. Here, $\Omega_{L,\infty}^{(\infty)}$ and Ω_L are the GNS vectors of $\omega_{L,\infty}^{(\infty)}$ and ω_L . Next, we specify W to be of the form $W = W_N(\xi)$ for $\xi \in \mathfrak{h}_{N,L}$ to write:

$$\begin{aligned} & \left\| \left[\pi_L(\beta_L(\alpha_{\infty}^{N'}(\eta_{L|t}^{(N')}(\alpha_{N'}^N(W)))) - \pi_L(\eta_{L|t}(\beta_L(\alpha_{\infty}^N(W)))) \right] \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L \right\| \\ &= \left\| \left[\pi_L(W_{\text{ct}}(R_{\infty}^{N'}(\tau_{L|t}^{(N')}(\alpha_{N'}^N(\xi)))) - \pi_L(W_{\text{ct}}(\tau_{L|t}(\alpha_{\infty}^N(\xi)))) \right] \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L \right\| \\ &= \left\| \left[\pi_L(W_{\text{ct}}(\xi_{N,N'}^{(t)})) - \pi_L(W_{\text{ct}}(\xi_{N,\infty}^{(t)})) \right] \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L \right\|, \end{aligned}$$

with the short hands $\xi_{N,N'}^{(t)}$ and $\xi_{N,\infty}^{(t)}$ for $R_{\infty}^{N'}(\tau_{L|t}^{(N')}(\alpha_{N'}^N(\xi)))$ and $\tau_{L|t}(\alpha_{\infty}^N(\xi))$ in the last line. By using the Weyl relations (3.1) and the explicit form (4.9) of the expectation values of Weyl operators in the state ω_L , we find (cp. [11, Proposition 5.2.29]):

$$\begin{aligned} & \left\| \left[\pi_L(W_{\text{ct}}(\xi_{N,N'}^{(t)})) - \pi_L(W_{\text{ct}}(\xi_{N,\infty}^{(t)})) \right] \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L \right\| \\ &\leq (\|R_{\infty}^M(\zeta)\|_{\mathfrak{h}_L} + \frac{1}{2}\|\xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L})\|\xi_{N,N'}^{(t)} - \xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L} + 2 \left(1 - e^{-\frac{1}{4}\|\xi_{N,N'}^{(t)} - \xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L}^2} \right) \\ &\leq (\|R_{\infty}^M(\zeta)\|_{\mathfrak{h}_L} + \frac{1}{2}\|\xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L} + \frac{1}{2})\|\xi_{N,N'}^{(t)} - \xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L} \\ &= (\|R_{\infty}^M(\zeta)\|_{\mathfrak{h}_L} + \frac{1}{2}\|R_{\infty}^N(\xi)\|_{\mathfrak{h}_L} + \frac{1}{2})\|\xi_{N,N'}^{(t)} - \xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L}, \end{aligned}$$

where we use the unitary implementation of τ_L on \mathfrak{h}_L . Thus, we obtain the anticipated Lipschitz estimate. Now, Proposition 4.8 states that for arbitrary $0 < T < \infty$:

$$\lim_{N' \rightarrow \infty} \sup_{|t| \leq T} \|\xi_{N,N'}^{(t)} - \xi_{N,\infty}^{(t)}\|_{\mathfrak{h}_L} = 0,$$

and, thus, we have,

$$\lim_{N' \rightarrow \infty} \sup_{|t| \leq T} \left\| \left(\pi_L(W_{\text{ct}}(\xi_{N,N'}^{(t)})) - \pi_L(W_{\text{ct}}(\xi_{N,\infty}^{(t)})) \right) \Psi \right\| = 0,$$

for the total set of vectors $\{\Psi \in \mathcal{H}_L \mid \Psi = \pi_L(W_{\text{ct}}(R_{\infty}^M(\zeta)))\Omega_L, \zeta \in \mathfrak{h}_{M,L}, M \in \mathbb{N}_0\}$. Finally, the result follows from combining this with the fact that the Weyl operators $W = W_N(\xi)$, $\xi \in \mathfrak{h}_{N,L}$, form a total set of uniformly bounded operators in $\mathcal{W}_{N,L}$. \square

Remark 4.10. We can identify \mathfrak{h}_L with a subspace of $\ell^2(\frac{\pi}{L}\mathbb{Z}^d, (2L)^{-d}\mu_{\frac{\pi}{L}\mathbb{Z}^d})^{\oplus 2}$ via the Fourier transform and the dispersion relation γ_m :

$$(\hat{f}, \hat{g}) \mapsto (\gamma_m^{-\frac{1}{2}}\hat{f}, \gamma_m^{\frac{1}{2}}\hat{g}).$$

This way τ_L is given by multiplication with the matrix-valued function $\exp(i\gamma_m\sigma_2)$, where σ_2 is the second Pauli matrix. Clearly, we can implement a dynamics given by γ_{μ_N} on $\ell^2(\frac{\pi}{L}\mathbb{Z}^d, (2L)^{-d}\mu_{\frac{\pi}{L}\mathbb{Z}^d})^{\oplus 2}$ using multiplication with $\exp(i\gamma_{\mu_N}\sigma_2)$. But, this is not the action of $\tau_L^{(N)}$ on the subspace $\mathfrak{h}_{N,L} \subset \mathfrak{h}_L$ and, thus, the convergence of the bounded matrix-functions

$$\exp(i\gamma_{\mu_N}\sigma_2) \rightarrow \exp(i\gamma_m\sigma_2), \quad N \rightarrow \infty,$$

is not the convergence of dynamics sought after.

By construction, $\eta_L : \mathbb{R} \curvearrowright \pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L})''$ equals the usual time evolution of the continuum free scalar field and it is implemented by a unitary group $U_{L|t} = e^{itH_L}$ with the (renormalized) free continuum Hamiltonian H_L as its generator. Explicitly, H_L is the second quantization of the generator h_L of τ_L on its natural domain [17]. Since γ_m is the free relativistic dispersion relation of mass m , we conclude that η_L has propagation speed $c = 1$, and we obtain a causal net of (spacetime) local von Neumann algebras for suitable $\mathcal{O} \subset \mathbb{R} \times \mathbb{T}_L^d$ [39]:

$$\mathcal{A}_L(\mathcal{O}) = \left(\bigcup_{t \in \mathbb{R}} \eta_{L|t}(\pi_{L,\infty}^{(\infty)}(\mathcal{W}_{\infty,L}(\mathcal{O}(t)))'' \right)'', \quad (4.28)$$

where $\mathcal{O}(t) = \{x \mid (t, x) \in \mathcal{O}\} \subset \mathbb{T}_L^d$.

Let us summarize our results on the scaling limit of the free ground states of the lattice scalar field by the following theorem:

Theorem 4.11. *Let $K \geq 6$. Let $m, \{\mu_N\}, \{\mathcal{W}_{N,L}\}, \{\alpha_{N'}^N\}, \{\omega_{L,M}^{(N)}\}$ as before. The scaling limit $\omega_{L,\infty}^{(\infty)}$ of $\{\omega_{L,M}^{(N)}\}$ gives rise to a continuum time-zero net of local algebras which is unitarily equivalent to the time-zero net of the free field with mass m on the torus \mathbb{T}_L^d .*

Remark 4.12. *In view of Figures 1 and 2, we note that while we used the scaling limit procedure to construct the horizontal sequences $\{\omega_{L,M}^{(N)}\}_{M \in \mathbb{N}_0}$ to arrive at the limit state $\omega_{L,\infty}^{(\infty)}$, we only considered the diagonal sequence of Hamiltonian $H_{L,0}^{(N)}$, or more precisely their associated unitary groups $\eta_L^{(N)}$, to obtain the Hamiltonian H_L associated with the continuum time evolution η_L .*

Remark 4.13. *There is yet another perspective on the convergence of dynamics because of the specific structure of the scaling maps α_{N+1}^N : At each finite scale N , the dynamics $\eta_L^{(N)}$ is unitarily implemented in the GNS representation of $\omega_{L,0}^{(N)}$, and because of the von Neumann uniqueness theorem there is an associated strongly-continuous, unitary one-parameter group $U^{(N)} : \mathbb{R} \rightarrow \mathcal{U}(\mathcal{H}_{N,L}^{(\infty)})$ on the GNS Hilbert space $\mathcal{H}_{N,L}^{(\infty)}$ of $\omega_{L,\infty}^{(N)}$. In Section 6.3 below, we show that the scaling maps α_{N+1}^N extend as normal, unital *-morphisms from $B(\mathcal{H}_{N,L})$ to $B(\mathcal{H}_{N+1,L})$, which in turn yields their extension from $B(\mathcal{H}_{N,L}^{(\infty)})$ to $B(\mathcal{H}_{N+1,L}^{(\infty)})$. This implies that each $U^{(N)}$ is represented in $\mathcal{U}(\mathcal{H}_{L,\infty}^{(\infty)})$ by¹³:*

$$U_{L|t}^{(N)} = \pi_{L,\infty}^{(\infty)}(\alpha_{\infty}^N(U_t^{(N)})), \quad t \in \mathbb{R}.$$

¹³The asymptotic unital *-morphisms $\alpha_{\infty}^N : B(\mathcal{H}_{N,L}) \rightarrow B_{\infty}$ is not necessarily normal, cf. Section 6.3 for the notation.

Therefore, as an alternative to the convergence considered in Corollary 4.9, we can analyze the convergence:

$$U_{L|t}^{(N)} \longrightarrow U_{L|t}, \quad N \longrightarrow \infty.$$

Locality by Lieb-Robinson bounds. Interestingly, there is another way to show a weaker form of locality via Lieb-Robinson bounds [65, 64]. To this end, we observe that the finite-scale dynamics $\eta_L^{(N)}$ extends to the C^* -inductive limit $\mathcal{W}_{\infty,L}$: For $N \in \mathbb{N}_0$ we consider the periodic extension of γ_{μ_N} from Γ_N to $\Gamma_\infty = \frac{\pi}{L}\mathbb{Z}^d$, which we also denote by γ_{μ_N} . Then, we extend $\tau_L^{(N)}$ to $\mathfrak{h}_{M,L}$ for all $M > N$ by

$$\tau_{L|t}^{(N)}(\hat{q}, \hat{p}) = (\cos(\gamma_{\mu_N} t)\hat{q} - \gamma_{\mu_N} \sin(\gamma_{\mu_N} t)\hat{p}, \quad \cos(\gamma_{\mu_N} t)\hat{p} + \gamma_{\mu_N}^{-1} \sin(\gamma_{\mu_N} t)\hat{q}), \quad (\hat{q}, \hat{p}) \in \mathfrak{h}_{M,L},$$

which is well-defined because the restriction of γ_{μ_N} to $\Gamma_M \subset \Gamma_\infty$ is well-defined by periodicity. Comparing the extension for $M' > M > N$, we find the consistency relation:

$$\tau_{L|t}^{(N)}(R_{M'}^M(\hat{q}, \hat{p})) = R_{M'}^M(\tau_{L|t}^{(N)}(\hat{q}, \hat{p})), \quad (\hat{q}, \hat{p}) \in \mathfrak{h}_{M,L}$$

which is due to the multiplicative character of $R_{M'}^M$ in momentum space, see (3.29). Therefore, we can extend $\tau_L^{(N)}$ to the inductive-limit symplectic space $\mathfrak{h}_{\infty,L}$ by:

$$\begin{aligned} \tau_{L|t}^{(N)}(R_\infty^M(\hat{q}, \hat{p})) &= \varepsilon_N^{\frac{d}{2}} \hat{\phi}(\varepsilon_N \cdot) (\cos(\gamma_{\mu_N} t)\hat{q} - \gamma_{\mu_N} \sin(\gamma_{\mu_N} t)\hat{p}, \cos(\gamma_{\mu_N} t)\hat{p} + \gamma_{\mu_N}^{-1} \sin(\gamma_{\mu_N} t)\hat{q}) \\ &= R_\infty^M(\cos(\gamma_{\mu_N} t)\hat{q} - \gamma_{\mu_N} \sin(\gamma_{\mu_N} t)\hat{p}, \cos(\gamma_{\mu_N} t)\hat{p} + \gamma_{\mu_N}^{-1} \sin(\gamma_{\mu_N} t)\hat{q}) \\ &= R_\infty^M(\tau_{L|t}^{(N)}(\hat{q}, \hat{p})), \end{aligned} \quad (4.29)$$

for any $M > N$.

This extension of $\tau_L^{(N)}$ to $\mathfrak{h}_{\infty,L}$ provides another justification for considering the convergence expressed in Proposition 4.8 because it implies that the convergence also holds on $\mathfrak{h}_{\infty,L} \subset \mathfrak{h}_L$ consistently over all scales. An explicit computation shows that the extension of $\tau_L^{(N)}$ is symplectic for each $M > N$, i.e. $\sigma_{M,L}(\tau_{L|t}^{(N)}(q, p), \tau_{L|t}^{(N)}(q', p')) = \sigma_{M,L}((q, p), (q', p'))$, which permits the extension of $\eta_L^{(N)}$ to $\mathcal{W}_{\infty,L}$ as a *-automorphism satisfying

$$\eta_{L|t}^{(N)}(\alpha_\infty^M(W_M(q, p))) = \alpha_\infty^M(\eta_{L|t}^{(N)}(W_M(q, p))) = \alpha_\infty^M(W_M(\tau_{L|t}^{(N)}(q, p))), \quad (4.30)$$

by general principles [26]. Now, we can invoke the Lieb-Robinson bounds for harmonic lattice systems [62] adapted to the Hamiltonian $H_{L,0}^{(N)}$ and the Weyl algebra $\mathcal{W}_{N,L}$:

Lemma 4.14. *For arbitrary subsets $X, Y \subset \Lambda_N$, all volume parameters $L = \varepsilon_N r_N > 0$, any one-particle vector $\xi, \xi' \in \mathfrak{h}_{N,L}$ with supports $\text{supp } \xi \subset X$, $\text{supp } \xi' \subset Y$, and any $\delta > 0$, we have:*

$$\left\| \left[\eta_{L|t}^{(N)}(W_N(\xi)), W_N(\xi') \right] \right\| \leq C_N \|\xi\|_{\infty,L} \|\xi'\|_{\infty,L} \sum_{x \in X, y \in Y} e^{-\frac{\delta}{\varepsilon_N} \left(d_N(x,y) - \frac{1}{2} c_{\mu_N} \max\left\{ \frac{2}{\delta}, e^{\frac{\delta}{2}+1} \right\} |t| \right)},$$

where

$$d_N(x, y) = \sum_{j=1}^d \min_{n_j \in \mathbb{Z}} |x_j - y_j + 2Ln_j|, \quad x, y \in \Lambda_N,$$

is the 1-distance on the torus \mathbb{T}_L^d restricted to Λ_N , and

$$c_{\mu_N} = (\mu_N^2 + 2d)^{\frac{1}{2}}, \quad C_N = 2 + c_{\mu_N} e^{\frac{\delta}{2}} + c_{\mu_N}^{-1}.$$

Proof. The estimate is a direct consequence of the estimate given in [62, Theorem 3.1] after reinstating the lattice spacing ε_N . \square

The estimate given in the Lemma leads to a speed of propagation $c' \geq 1$:

Proposition 4.15. *Let $K \geq 2$. Provided the renormalization condition (4.18) holds, there exists a constant $1 \leq c' \leq d^{\frac{1}{2}} \max\{\frac{2}{\delta}, e^{\frac{\delta}{2}+1}\}$, the scaling-limit Lieb-Robinson velocity, such that for $S' \cap S_{c'T} = \emptyset$ with $S_{c'T} = \{x \mid \text{dist}(x, S) < c'T\}$,*

$$\lim_{N \rightarrow \infty} \left\| \left[\eta_{L|t}^{(N)}(W), W' \right] \right\| = 0,$$

for all $W \in \mathcal{W}_{\infty, L}(S)$, $W' \in \mathcal{W}_{\infty, L}(S')$ exponentially fast and uniformly for $|t| < T$.

Proof. First, we consider two elementary Weyl operators $W, W' \in \mathcal{W}_{\infty, L}$. By construction there exist $N, N' \in \mathbb{N}_0$ and $\xi \in \mathfrak{h}_{N, L}$, $\xi' \in \mathfrak{h}_{N', L}$ such that $W = \alpha_{\infty}^N(W_N(\xi))$ and $W' = \alpha_{\infty}^{N'}(W_{N'}(\xi'))$ (without loss of generality we may assume $N \leq N'$). For any $M \geq N' \geq N$ we have:

$$\left[\eta_{L|t}^{(M)}(W), W' \right] = \alpha_{\infty}^M \left(\left[\eta_{L|t}^{(M)}(W_M(R_M^N(\xi))), W_M(R_M^{N'}(\xi')) \right] \right).$$

Since α_{∞}^M is an injective C^* -homomorphism, we can apply the estimate of Lemma 4.14:

$$\left\| \left[\eta_{L|t}^{(M)}(W), W' \right] \right\| \leq C_M \|R_M^N(\xi)\|_{\infty, L} \|R_M^{N'}(\xi')\|_{\infty, L} \sum_{x \in X, y \in Y} e^{-\frac{\delta}{\varepsilon_M} \left(d_M(x, y) - \frac{1}{2} c_{\mu_M} \max\left\{ \frac{2}{\delta}, e^{\frac{\delta}{2}+1} \right\} |t| \right)},$$

and by the renormalization conditions, we have

$$\lim_{M \rightarrow \infty} c_{\mu_M} = 2d^{\frac{1}{2}}, \quad \lim_{M \rightarrow \infty} C_M = 2 + 2d^{\frac{1}{2}} e^{\frac{\delta}{2}} + 2^{-1} d^{-\frac{1}{2}} = C_d.$$

Now, if $\text{supp } R_{\infty}^N(q_{\xi}, p_{\xi}) \subset S$ and $\text{supp } R_{\infty}^{N'}(q_{\xi'}, p_{\xi'}) \subset S'$, we can apply the estimate with $\text{dist}_1(S, S') \leq \lim_{M \rightarrow \infty} d_M(x, y) < \infty$, where $\text{dist}_1(S, S')$ is the 1-distance between subsets of \mathbb{T}_L^d . Moreover, by (3.7),

$$\begin{aligned} \|R_M^N(\xi)\|_{\infty, L} &= \varepsilon_M^{\frac{d}{2}} \left\| \varepsilon_M^{\frac{1}{2}} R_M^N(q_{\xi}) + i \varepsilon_M^{-\frac{1}{2}} R_M^N(p_{\xi}) \right\|_{\infty, L}, \\ \|R_M^{N'}(\xi')\|_{\infty, L} &= \varepsilon_M^{\frac{d}{2}} \left\| \varepsilon_M^{\frac{1}{2}} R_M^{N'}(q_{\xi'}) + i \varepsilon_M^{-\frac{1}{2}} R_M^{N'}(p_{\xi'}) \right\|_{\infty, L}, \end{aligned}$$

which can be combined with (4.21) & (4.22) for $\alpha = 0$ in Lemma 4.2 to show:

$$\lim_{M \rightarrow \infty} \|R_M^N(q_{\xi})\|_{\infty, L} \leq \|\hat{\phi}^{(\varepsilon_N)} \hat{q}_{\xi}\|_{L^1, L} < \infty, \quad \lim_{M \rightarrow \infty} \|R_M^{N'}(p_{\xi'})\|_{\infty, L} \leq \|\hat{\phi}^{(\varepsilon_N)} \hat{p}_{\xi'}\|_{L^1, L} < \infty,$$

for all $N \in \mathbb{N}_0$, and similarly for ξ' , where $\|\cdot\|_{L^1, L}$ is the norm of $L^1(\frac{\pi}{L} \mathbb{Z}^d, (2L)^{-d} \mu_{\frac{\pi}{L} \mathbb{Z}^d})$. Thus, we find (in the limit $M \rightarrow \infty$):

$$\begin{aligned} \left\| \left[\eta_{L|t}^{(M)}(W), W' \right] \right\| &\lesssim C_d \varepsilon_M^{-d} \left(\varepsilon_M^{\frac{1}{2}} \|\hat{\phi}^{(\varepsilon_N)} \hat{q}_{\xi}\|_{L^1, L} + \varepsilon_M^{-\frac{1}{2}} \|\hat{\phi}^{(\varepsilon_N)} \hat{p}_{\xi}\|_{L^1, L} \right) \\ &\quad \times \left(\varepsilon_M^{\frac{1}{2}} \|\hat{\phi}^{(\varepsilon_N)} \hat{q}_{\xi'}\|_{L^1, L} + \varepsilon_M^{-\frac{1}{2}} \|\hat{\phi}^{(\varepsilon_N)} \hat{p}_{\xi'}\|_{L^1, L} \right) \\ &\quad \times \text{vol}(S) \text{vol}(S') e^{-\frac{\delta}{\varepsilon_M} \left(\text{dist}_1(S, S') - d^{\frac{1}{2}} \max\left\{ \frac{2}{\delta}, e^{\frac{\delta}{2}+1} \right\} |t| \right)}. \\ &\rightarrow 0, \end{aligned}$$

where the expressions $\text{vol}(S)$, $\text{vol}(S')$ arise by collecting an overall factor ε_M^{2d} in front of the sum $\sum_{x \in X, y \in Y}$ and choosing $X = \Lambda_M \cap S$, $Y = \Lambda_M \cap S'$:

$$\text{vol}(S) = \lim_{M \rightarrow \infty} \varepsilon_M^d \sum_{x \in \Lambda_M \cap S} 1, \quad \text{vol}(S') = \lim_{M \rightarrow \infty} \varepsilon_M^d \sum_{y \in \Lambda_M \cap S'} 1.$$

Using the equivalence of dist_1 and dist , i.e. $\text{dist} \leq \text{dist}_1 \leq d^{\frac{1}{2}} \text{dist}$, and $\lim_{M \rightarrow \infty} \varepsilon_M^{-k} e^{-\frac{\delta}{\varepsilon_M}} = 0$ for any $k \in \mathbb{Z}$, the statement is proved for elementary Weyl operators and their finite linear combinations with the scaling-limit Lieb-Robinson velocity bounded by:

$$1 \leq \max \left\{ \frac{2}{\delta}, e^{\frac{\delta}{2}+1} \right\} \leq c' \leq d^{\frac{1}{2}} \max \left\{ \frac{2}{\delta}, e^{\frac{\delta}{2}+1} \right\}.$$

Then, because $\eta_L^{(M)}$ is a group of *-automorphisms of $\mathcal{W}_{\infty, L}$ and Weyl operators form a bounded total set in $\mathcal{W}_{\infty, L}(S)$, $\mathcal{W}_{\infty, L}(S')$, we obtain the desired statement. \square

Corollary 4.16. *Let $K \geq 2$. With the scaling-limit Lieb-Robinson velocity c' from Proposition 4.15 and $S' \cap S_{c'T} = \emptyset$ with $S_{c'T} = \{x \mid \text{dist}(x, S) < c'T\}$,*

$$\text{SOT-lim}_{N \rightarrow \infty} \left[\eta_{L|t}^{(N)}(W), W' \right] = 0,$$

for all $W \in \mathcal{W}_{\infty, L}(S)$, $W' \in \mathcal{W}_{\infty, L}(S')$ exponentially fast and uniformly for $|t| < T$ relative to the representation given by $\omega_{L, \infty}^{(\infty)}$. Moreover, for all such $t \in \mathbb{R}$ and for all $W \in \pi_{L, \infty}^{(\infty)}(\mathcal{W}_{\infty, L}(S))''$, $W' \in \pi_{L, \infty}^{(\infty)}(\mathcal{W}_{\infty, L}(S'))''$:

$$[\eta_{L|t}(W), W'] = 0.$$

Proof. The statement about SOT (strong operator topology) convergence is immediate from the norm convergence of Proposition 4.15. The commutativity for weak closures follows from the the SOT convergence of $\eta_L^{(N)}$ to η_L and the SOT continuity of η_L . \square

Remark 4.17. *Note, that even in the case $\text{dist}_1 = d^{\frac{1}{2}} \text{dist}$, the (optimal) scaling-limit Lieb-Robinson velocity is $c' = \frac{2}{\delta_0} \sim 3.59$, given by the Lambert W function,*

$$\frac{\delta_0}{2} e^{\frac{\delta_0}{2}} = e^{-1},$$

which is still far from the correct propagation speed $c = 1$.

4.2.5 Infinite volume limit

Here we compare the finite volume theory and the infinite volume theory. Given a bounded open set $S \subset \mathbb{R}^d$, one has $S \Subset (-L, L)^d$ for sufficiently large L . The C^* -algebra $\mathcal{W}_{\infty, L}(S)$ actually does not depend on L (possibly $L = \infty$). Indeed, with the notation $\mathfrak{h}_{N, L}(S) := \ell^2(\Lambda_N(S))$ (cf. Sec. 2.3), $\beta_L(\mathcal{W}_{\infty, L}(S)) \subset \mathcal{W}(\mathfrak{h}_L)$ is generated by Weyl operators $W_{\text{ct}}(\xi)$, $\xi \in \mathfrak{h}_{\infty, L}(S) := \bigcup_N R_{\infty}^N \mathfrak{h}_{N, L}(S)$, and each $\xi \in \mathfrak{h}_{\infty, L}(S)$ can be also thought as a compactly supported function on \mathbb{R}^d , and $\sigma_L(\xi, \eta) = \sigma_{\infty}(\xi, \eta)$ for $\xi, \eta \in \mathfrak{h}_L(S)$ (see (4.8), (4.14)). Therefore, $\mathcal{W}_{\infty, L}(S)$ can also be identified with the C^* -subalgebra of the infinite volume free field C^* -algebra $\mathcal{W}(\mathfrak{h}_{\infty})$ generated by the Weyl operators $W_{\text{ct}}(\xi)$, $\xi \in \mathfrak{h}_{\infty, L}(S)$.

With this embedding and Lemma 4.3 in mind, we can consider on $\mathcal{W}_{\infty, L}(S)$ the family of continuum states $\{\omega_L\}_{L>0}$, and the limit $L \rightarrow \infty$.

Proposition 4.18. *For each $\xi \in \mathfrak{h}_{\infty,L}(S)$ we have*

$$\lim_{L \rightarrow +\infty} \omega_L(W_{\text{ct}}(\xi)) = \omega_{\infty}(W_{\text{ct}}(\xi)),$$

and $\{\omega_L\}$ converges to ω_{∞} in the weak* topology on $\mathcal{W}_{\infty,L}(S)$.

Proof. It is enough to compute the exponent which appears in the expectation values:

$$\lim_{L \rightarrow +\infty} \frac{1}{(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \left| \frac{\hat{q}_{\xi}(k)}{\gamma_m^{\frac{1}{2}}(k)} + i\gamma_m^{\frac{1}{2}}(k)\hat{p}_{\xi}(k) \right|^2 = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} d^d k \left| \frac{\hat{q}_{\xi}(k)}{\gamma_m^{\frac{1}{2}}(k)} + i\gamma_m^{\frac{1}{2}}(k)\hat{p}_{\xi}(k) \right|^2.$$

Indeed, if the right-hand side is integrable, one can restrict it to a sufficiently large compact interval with a small error, and the corresponding restriction of the sum on the left hand side is then a corresponding Riemann sum, which is convergent in the $L \rightarrow \infty$ limit due to the regularity of $\hat{q}_{\xi}, \hat{p}_{\xi}$. The weak* convergence follows by approximation in norm. \square

We now prove that the GNS representations with respect to the vacuum states with different volumes L are locally equivalent. This enables us to identify locally the finite volume algebras in a cylinder and in the Minkowski space, showing that we can obtain the local algebras of the infinite volume continuum free field through our finite volume scaling limit procedure.

Following the steps of [20, Appendix A], we first show that the two representations are quasi-equivalent by the main theorem in [3], then this is actually unitary equivalence because local algebras are type III factors. The assumptions of [3] are satisfied if one proves the following facts.

1. The topologies induced by the two-point functions on the one-particle spaces are equivalent.
2. The kernels of the quadratic forms associated to the difference of the two point functions on the torus and on \mathbb{R}^d (Q_+ and Q_- in Lemma 4.21) are sufficiently regular. This entails the Hilbert-Schmidt property of the Araki-Yamagami operator.

We show them in Lemma 4.20 and Lemma 4.21, respectively. As a preparation for the equivalence of topologies we first prove two versions of the Young inequality relating continuous L^2 -norms with discrete convolutions and vice versa.

Lemma 4.19. *Given $f \in \mathcal{S}(\mathbb{R})$, $g \in L^2(\mathbb{R})$, there holds*

$$\begin{aligned} \int_{\mathbb{R}} dk \left| \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)||g(p)| \right|^2 &\leq \max_{k \in [0, \frac{\pi}{L}]} \|f_k\|_{\ell^1(\frac{\pi}{L}\mathbb{Z})} \|f\|_{L^1(\mathbb{R})} \|g\|_{\ell^2(\frac{\pi}{L}\mathbb{Z})}^2, \\ \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \left| \int_{\mathbb{R}} dp |f(k-p)||g(p)| \right|^2 &\leq \max_{k \in [0, \frac{\pi}{L}]} \|f_k\|_{\ell^1(\frac{\pi}{L}\mathbb{Z})} \|f\|_{L^1(\mathbb{R})} \|g\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

where $f_k(p) := f(k-p)$, $k, p \in \mathbb{R}$ and $\|h\|_{\ell^1(\frac{\pi}{L}\mathbb{Z})} = \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |h(k-p)|$.

Proof. There holds

$$\begin{aligned} \int_{\mathbb{R}} dk \left| \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)||g(p)| \right|^2 &= \int_{\mathbb{R}} dk \left| \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)|^{1/2} |g(p)| |f(k-p)|^{1/2} \right|^2 \\ &\leq \int_{\mathbb{R}} dk \left[\sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)||g(p)|^2 \right] \left[\sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)| \right], \end{aligned}$$

as the last expression contains more positive terms than the previous one. Since $f \in \mathcal{S}(\mathbb{R})$, the function $k \mapsto \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)| = \|f_k\|_{\ell^1(\frac{\pi}{L}\mathbb{Z})}$ is $\frac{\pi}{L}$ -periodic and continuous (the series converges uniformly in $k \in [0, \frac{\pi}{L}]$), and therefore $\sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)| \leq \max_{k \in [0, \frac{\pi}{L}]} \|f_k\|_{\ell^1(\frac{\pi}{L}\mathbb{Z})} < \infty$ for all $k \in \mathbb{R}$. Moreover

$$\int_{\mathbb{R}} dk \left[\sum_{p \in \frac{\pi}{L}\mathbb{Z}} |f(k-p)| |g(p)|^2 \right] = \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |g(p)|^2 \int_{\mathbb{R}} dk |f(k-p)| = \|g\|_{\ell^2(\frac{\pi}{L}\mathbb{Z})}^2 \|f\|_{L^1(\mathbb{R})},$$

which proves the first inequality. The second one is proven in the same way, exchanging the roles of sums and integrals. \square

Let us show the first of the conditions for unitary equivalence.

Lemma 4.20. *For $S \Subset (-L, L)$ the norms $\|\cdot\|_L$ and $\|\cdot\|_{\infty}$ defined by (4.7) and (4.13) respectively, are equivalent on $\mathfrak{h}_{\infty, L}(S)$.*

Proof. Setting, for a function $\xi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\|\xi\|_{L, \pm}^2 := \frac{1}{2L} \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \gamma_m(k)^{\pm 1} |\hat{\xi}(k)|^2, \quad \|\xi\|_{\infty, \pm}^2 := \frac{1}{2\pi} \int_{\mathbb{R}} dk \gamma_m(k)^{\pm 1} |\hat{\xi}(k)|^2,$$

it is sufficient to show that on $\mathfrak{h}_{\infty, L}(S)$ the norm $\|\cdot\|_{L, \pm}$ is equivalent to $\|\cdot\|_{\pm}$ respectively.

To this end, let $\xi \in \mathfrak{h}_{\infty, L}(S)$ be a real function, and let $\chi \in C_c^{\infty}((-L, L))$ be such that $\chi(x) = 1$ for all $x \in \text{supp } \xi$. Since $\text{supp } \xi \subset (-L, L)$, we can define its $2L$ -periodization

$$P\xi(x) := \sum_{n \in \mathbb{Z}} \xi(x + 2Ln) = \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \hat{\xi}(k) e^{ikx},$$

and thanks to $\sum_{k \in \frac{\pi}{L}\mathbb{Z}} |\hat{\xi}(k)| < +\infty$ we obtain, for all $k \in \mathbb{R}$,

$$\begin{aligned} \hat{\xi}(k) &= \widehat{\chi P\xi}(k) = \int_{\mathbb{R}} dx \chi(x) P\xi(x) e^{-ikx} = \sum_{p \in \frac{\pi}{L}\mathbb{Z}} \hat{\xi}(p) \int_{\mathbb{R}} dx \chi(x) e^{-i(k-p)x} \\ &= \sum_{p \in \frac{\pi}{L}\mathbb{Z}} \hat{\chi}(k-p) \hat{\xi}(p). \end{aligned}$$

Now since $k \mapsto \gamma_m(k)$ is a convex function,

$$\gamma_m(k) = \gamma_m(k-p+p) \leq \frac{1}{2}(\gamma_m(2(k-p)) + \gamma_m(2p)) = \gamma_{\frac{m}{2}}(k-p) + \gamma_{\frac{m}{2}}(p) \leq \gamma_m(k-p) + \gamma_m(p),$$

which entails $\gamma_m(k)^{\frac{1}{2}} \leq \gamma_m(k-p)^{\frac{1}{2}} + \gamma_m(p)^{\frac{1}{2}}$. As a consequence

$$\gamma_m(k)^{\frac{1}{2}} |\hat{\xi}(k)| \leq \sum_{p \in \frac{\pi}{L}\mathbb{Z}} \gamma_m(k-p)^{\frac{1}{2}} |\hat{\chi}(k-p)| |\hat{\xi}(p)| + \sum_{p \in \frac{\pi}{L}\mathbb{Z}} |\hat{\chi}(k-p)| \gamma_m(p)^{\frac{1}{2}} |\hat{\xi}(p)|,$$

and in view of the fact that $\hat{\chi}, \gamma_m^{\frac{1}{2}} \hat{\chi} \in \mathcal{S}(\mathbb{R})$, from the first inequality of Lemma 4.19 the (squared) $L^2(\mathbb{R})$ -norms of the two terms in the right hand side can be estimated by $C \|\hat{\xi}\|_{\ell^2(\frac{\pi}{L}\mathbb{Z})}^2 \leq 2LC \|\xi\|_{L, +}^2$ and by $C \|\gamma_m^{\frac{1}{2}} \hat{\xi}\|_{\ell^2(\frac{\pi}{L}\mathbb{Z})}^2 = 2LC \|\xi\|_{L, +}^2$ respectively, for a suitable constant $C > 0$. This implies

$$\|\xi\|_{\infty, +}^2 = \frac{1}{2\pi} \|\gamma_m^{1/2} \hat{\xi}\|_{L^2(\mathbb{R})}^2 \leq \frac{4LC}{\pi} \|\xi\|_{L, +}^2.$$

Conversely, we can also write, for all $k \in \frac{\pi}{L}\mathbb{Z}$,

$$\begin{aligned} \gamma_m(k)^{\frac{1}{2}}|\hat{\xi}(k)| &= \gamma_m(k)^{\frac{1}{2}}|\widehat{\chi\xi}(k)| \leq \gamma_m(k)^{\frac{1}{2}} \int_{\mathbb{R}} dp |\hat{\chi}(k-p)| |\hat{\xi}(p)| \\ &\leq \int_{\mathbb{R}} dp \gamma_m(k-p)^{\frac{1}{2}} |\hat{\chi}(k-p)| |\hat{\xi}(p)| + \int_{\mathbb{R}} dp |\hat{\chi}(k-p)| \gamma_m(p)^{\frac{1}{2}} |\hat{\xi}(p)|, \end{aligned}$$

and using now the second inequality of the previous Lemma we obtain, arguing as above,

$$\|\xi\|_{L,+}^2 = \frac{1}{2L} \|\gamma_m^{1/2} \hat{\xi}\|_{\ell^2(\frac{\pi}{L}\mathbb{Z})}^2 \leq \frac{4\pi C}{L} \|\xi\|_{\infty,+}^2,$$

which shows the required equivalence of $\|\cdot\|_{L,+}$ and $\|\cdot\|_{\infty,+}$.

In order to prove the equivalence of the norms $\|\cdot\|_{L,-}$, $\|\cdot\|_{\infty,-}$, we start by observing that

$$\begin{aligned} \frac{\gamma_m(p)^{1/2}}{\gamma_m(k)^{1/2}} &\leq C \left[\frac{1+|p|^2}{1+|k|^2} \right]^{1/4} \leq C' \left[\frac{1+|p|}{1+|k|} \right]^{1/2} \leq C' \left[\frac{1+|k|+|p-k|}{1+|k|} \right]^{1/2} \\ &\leq C'(1+|k-p|)^{1/2}, \end{aligned}$$

for suitable constants $C, C' > 0$, where in the second inequality we used the fact that the function $k \mapsto (1+|k|^2)^{1/2}/(1+|k|)$ is bounded, continuous, non-vanishing on \mathbb{R} and converges to 1 for $|k| \rightarrow +\infty$. We obtain then as above the inequality

$$\frac{1}{\gamma_m(k)^{1/2}} |\hat{\xi}(k)| \leq C' \sum_{p \in \frac{\pi}{L}\mathbb{Z}} (1+|k-p|)^{1/2} |\hat{\chi}(k-p)| \frac{1}{\gamma_m(p)^{1/2}} |\hat{\xi}(p)|,$$

and the analogous one with the integral over $p \in \mathbb{R}$ replacing the sum. Using then the fact that $k \mapsto (1+|k|)^{1/2} |\hat{\chi}(k)|$ is in $\mathcal{S}(\mathbb{R})$ the equivalence of the norms is proven by the same argument based on the previous lemma as above. \square

Lemma 4.21. *For $S \Subset (-L, L)$ there are smooth functions Q_{\pm} on $(-2L, 2L)$ such that, for all real functions $\xi, \eta \in \mathfrak{h}_{\infty, L}(S)$*

$$\int_{\mathbb{R}} dk \frac{1}{\gamma_m(k)} \overline{\hat{\xi}(k)} \hat{\eta}(k) - \frac{\pi}{L} \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \frac{1}{\gamma_m(k)} \overline{\hat{\xi}(k)} \hat{\eta}(k) = \int_{S \times S} dx dy Q_-(x-y) \xi(x) \eta(y), \quad (4.31)$$

$$\int_{\mathbb{R}} dk \gamma_m(k) \overline{\hat{\xi}(k)} \hat{\eta}(k) - \frac{\pi}{L} \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \gamma_m(k) \overline{\hat{\xi}(k)} \hat{\eta}(k) = \int_{S \times S} dx dy Q_+(x-y) \xi(x) \eta(y). \quad (4.32)$$

Proof. By [1, Formula 9.6.21]

$$\int_{\mathbb{R}} dk \frac{1}{\gamma_m(k)} \overline{\hat{\xi}(k)} \hat{\eta}(k) = 2 \int_{S \times S} dx dy K_0(m|x-y|) \xi(x) \eta(y)$$

where $K_0(z)$, $z > 0$, is the modified Bessel function of order zero, which is positive, smooth for $z \neq 0$ and satisfies

$$K_0(z) \leq e^{-z}$$

for z sufficiently large. Moreover, the Poisson summation formula claims that

$$\frac{\pi}{L} \sum_{k \in \mathbb{Z}} \delta \left(x - k \frac{\pi}{L} \right) = \sum_{k \in \mathbb{Z}} e^{i2Lkx}$$

in the distributional sense, and therefore, since $k \mapsto \frac{1}{\gamma_m(k)} \widehat{\xi}(k) \hat{\eta}(k)$ decreases fast enough for $|k| \rightarrow +\infty$ [72, Corollary VII. 2.6],

$$\begin{aligned} \frac{\pi}{L} \sum_{k \in \frac{\pi}{L}\mathbb{Z}} \frac{1}{\gamma_m(k)} \widehat{\xi}(k) \hat{\eta}(k) &= \sum_{n \in \mathbb{Z}} \int_{\mathbb{R}} dk \frac{1}{\gamma_m(k)} \widehat{\xi}(k) \hat{\eta}(k) e^{-i2Lnk} \\ &= 2 \int_{S \times S} dx dy \left[\sum_{n \in \mathbb{Z}} K_0(m|x - y - 2Ln|) \right] \xi(x) \eta(y) \end{aligned}$$

where the sum inside the integral converges uniformly on $S \times S$ thanks to the exponential decay of K_0 . Therefore, looking at (4.31) we find the following expression for Q_-

$$Q_-(x) = \sum_{n \in \mathbb{Z} \setminus \{0\}} 2 K_0(m|x - 2Ln|)$$

The series is term by term smooth in $(-2L, 2L)$ since it does not contain the term with $n = 0$. Moreover, one can see that the s -th derivative of K_0 can be written in terms of K_0 and K'_0 , by the following formulas (cf. [1, sec. 9.6])

$$K'_0(z) = -K_1(z), \quad z^2 K''_0(z) + z K_0(z)' - z^2 K_0(z) = 0$$

and one gets therefore

$$K_0^{(s)}(z) = P_{s-2}(1/z) K_0(z) + R_{s-1}(1/z) K'_0(z), \quad s \geq 2,$$

where P_s, R_s are polynomials. As a consequence, since also $|K_1(z)| \leq e^{-z}$ for $z \rightarrow +\infty$, $K_0^{(s)}$ is exponentially decaying for $z \rightarrow +\infty$ too, and therefore the term by term s -th derivative of the series defining Q_- is uniformly convergent on $S \times S$, showing that Q_- is smooth on $(-2L, 2L)$.

Concerning (4.32), by the argument in the proof of [20, Lemma A.5], there holds

$$\int_{\mathbb{R}} dk \gamma_m(k) \widehat{\xi}(k) \hat{\eta}(k) = - \int_{S \times S} dx dy \frac{m}{|x - y|} K_1(m|x - y|) \xi(x) \eta(y),$$

and therefore, by a similar argument as the above one,

$$Q_+(x) = - \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{m}{|x - 2Ln|} K_1(m|x - 2Ln|)$$

is smooth in $(-2L, 2L)$. □

Theorem 4.22. *Let $K \geq 6$. For each bounded open set $S \Subset (-L, L)^d$, the finite volume representation $\pi_{L, \infty}^{(\infty)} \cong \pi_L$ of $\mathcal{W}_{\infty, L}(S)$ is unitarily equivalent to the infinite volume representation π_{∞} .*

Proof. Assume $d = 1$. Consider the completion of the space of real functions in $\mathfrak{h}_L(S)$ (thought as compactly supported functions on \mathbb{R}) with respect to the norms $\|\cdot\|_{\pm}$, and, on these spaces, the operators Q_{\pm} defined by

$$\langle \xi, Q_{\pm} \eta \rangle_{\pm} = \int_{S \times S} dx dy Q_{\pm}(x-y) \xi(x) \eta(y).$$

Then using Lemma 4.21 and the argument of the proofs of Lemmas A.4 and A.6 of [20], one verifies that Q_{\pm} are trace-class, and this, together with Lemma 4.20, implies the quasiequivalence statement by [3].

Now, as we take $K \geq 6$ and the density result (Lemma 4.6) is a local property and hence holds also for $L = \infty$, we have $\pi_{\infty}(\mathcal{W}_{\infty, L}(S))'' = \pi_{\infty}(\mathcal{W}_L(S))''$, and the latter is a type III factor [2]. Then also $\pi_{L, \infty}^{(\infty)}(\mathcal{W}_{\infty, L}(S))''$ is a type III factor. Therefore the last statement follows directly from [24, Sect. 5.6.6].

If $d > 1$, Lemmas 4.19 and 4.20 and their proofs continue to hold with the obvious modifications. Concerning lemma 4.21, the smoothness of the kernels Q_{\pm} around $x = 0$ can be proven by [68, IX.46], providing the smoothness of the infinite volume 2-point function in spatial directions away from the origin, and by a Payley-Wiener argument, providing its exponential decay. \square

Remark 4.23. *The spatial isomorphism defined in Theorem 4.22 depends in principle on S and L with $S \Subset (-L, L)^d$, and should be denoted by $\theta_{S, L}$. However for $\xi \in \mathfrak{h}_L(S)$ there holds*

$$\theta_{S, L}(\pi_L(W_{\text{ct}}(\xi))) = \pi_{\infty}(W_{\text{ct}}(\xi)),$$

where on the left side ξ is considered as function on \mathbb{T}_L^d and on the right side as a compactly supported function on \mathbb{R}^d . As a consequence, given tori \mathbb{T}_L^d and $\mathbb{T}_{L'}^d$, with $L < L'$ and $S \Subset (-L, L)^d$, then

$$\theta_{S, L}(\pi_L(\mathcal{W}_L(S))'') = \theta_{S, L'}(\pi_{L'}(\mathcal{W}_{L'}(S))'').$$

Similarly, if $S \subset S' \Subset (-L, L)^d$ then the consistency property $\theta_{S', L} \upharpoonright \pi_L(\mathcal{W}_L(S))'' = \theta_{S, L}$ holds. Because of this, we will use the simplified notation θ when no confusion can arise.

In view of the above remark, it also follows that θ locally intertwines the action of space translations on the torus and on \mathbb{R}^d , namely

$$\theta \text{Ad} U_L(a, 0) \upharpoonright \mathcal{A}_L(S) = \text{Ad} U_{\infty}(a, 0) \theta \upharpoonright \mathcal{A}_L(S),$$

where $S \subset \mathbb{R}^d$ is a bounded open set, $a \in \mathbb{R}^d$ and L is large enough that the closures of S and $S + a$ are both contained in $(-L, L)^d$.

A similar statement also holds for time translations. Indeed, in view of the fact that the solutions of the Klein-Gordon equation of the torus have propagations speed $c = 1$, with $\mathcal{B}_1(0)$ the unit ball centered at the origin, there holds

$$\text{Ad} U_L(0, t)(\mathcal{A}_L(S)) = \mathcal{A}_L(S + |t|\mathcal{B}_1(0))$$

whenever S is a bounded open set, $t \in \mathbb{R}$ and L is large enough that S and $S + t\mathcal{B}_1(0)$ are both contained in $(-L, L)^d$. This clearly entails

$$\theta \text{Ad} U_L(0, t) \upharpoonright \mathcal{A}_L(S) = \text{Ad} U_{\infty}(0, t) \theta \upharpoonright \mathcal{A}_L(S).$$

5 Other scaling-map schemes

From the perspective of the wavelet scaling map of Section 3.3 being a generalization of the block-spin scaling map, we discuss the relations between other scaling map schemes in this section: First, the block-spin case again. Second, a singular real-space scheme leading to point-like localization of lattice operators in the scaling limit in Section 5.2. Third, a momentum-space scheme implementing sharp momentum cutoff in Section 5.3.

The last two of these schemes can be formulated analogous to the wavelet scaling map at the one-particle level as in (3.25) in Definition 3.3. The main difference consists in the use of low-pass filters $\{h_n\}_{n \in \mathbb{Z}^d}$ concentrated at $n = 0$ (point-like localization) or scale-dependent full support, $h_n = h_n^{(N)}$, $\text{supp } h^{(N)} = \mathbb{Z}_{2r_N}^d$ (sharp momentum cutoff). In contrast with the wavelet scaling map this results in the drawback that these schemes do not directly lead to local algebras in terms of lattice operators in scaling limit. More specifically, the block spin renormalization does not have enough regularity to include the momentum operator, point-like localization leads to sharply localized field-like objects and the scaling limit of initial states can only be defined in a distributional sense necessitating a Wightman reconstruction, while sharp momentum cutoff leads to a scaling limit in terms of fully non-local algebras also requiring an indirect reconstruction of local operators.

5.1 The block-spin transformation case

The expression (4.19) does not define limit states $\omega_{L,\infty}^{(N)}$ on $\mathcal{W}_{N,L}$ if the renormalization group define by the block-spin scaling map (3.12). Nevertheless, it is still possible to interpret (4.19) in a distributional sense. To this end, we exploit the fact that the states $\omega_{\mu_{N+M},M} = \omega_{L,M}^{(N)}$ are quasi-free and, thus, are determined by there two-point functions:

$$\begin{aligned} W_{L,M|\Phi\Phi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{-\frac{1+d}{2}} \Phi_N(x) \varepsilon_N^{-\frac{1+d}{2}} \Phi_N(y))) \\ &= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(t\delta_x^{(N)})) \alpha_{N+M}^N(W_N(s\delta_y^{(N)})))|_{t=s=0} \\ &= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N^{-1} \gamma_{\mu_{N+M}}(k)^{-1} \\ &\quad \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(x' - x) \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(y' - y) e^{ik \cdot (x' - y')}, \end{aligned}$$

$$\begin{aligned} W_{L,M|\Pi\Pi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{\frac{1-d}{2}} \Pi_N(x) \varepsilon_N^{\frac{1-d}{2}} \Pi_N(y))) \\ &= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(it\delta_x^{(N)})) \alpha_{N+M}^N(W_N(is\delta_y^{(N)})))|_{t=s=0} \\ &= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N \gamma_{\mu_{N+M}}(k) \\ &\quad \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(x' - x) \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(y' - y) e^{ik \cdot (x' - y')}, \end{aligned}$$

$$\begin{aligned} W_{L,M|\Phi\Pi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{-\frac{1+d}{2}} \Phi_N(x) \varepsilon_N^{\frac{1-d}{2}} \Pi_N(y))) \\ &= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(t\delta_x^{(N)})) \alpha_{N+M}^N(W_N(is\delta_y^{(N)})))|_{t=s=0} \\ &= \frac{i}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(x' - x) \varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}(y' - y) e^{ik \cdot (x' - y')}, \end{aligned}$$

But, these expression are well-defined for arbitrary $x, y \in \mathbb{T}_L^d$ and consistent with respect to the inclusions $\Lambda_N \subset \Lambda_{N'} \subset \mathbb{T}_L^d$ for $N < N'$ as a consequence of the scaling equation (3.19).

Then, as a consequence of the Paley-Wiener-Schwartz characterization of the (continuum) Fourier transform of a functions $f, f' \in C^\infty(\mathbb{T}_L^d)$, the expression,

$$\begin{aligned} W_{L,M|\Phi\Phi}^{(N)}(f, f') &= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N^{-1} \gamma_{\mu_{N+M}}(k)^{-1} \\ &\quad \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}, \\ W_{L,M|\text{III}}^{(N)}(f, f') &= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N \gamma_{\mu_{N+M}}(k) \\ &\quad \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}, \\ W_{L,M|\Phi\Pi}^{(N)}(f, f') &= \frac{i}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_{N+M}^{2d} \sum_{x', y' \in \Lambda_{N+M}} \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}, \end{aligned}$$

admit well-defined limits for $M \rightarrow \infty$ (by a Riemann-sum argument):

$$\begin{aligned} W_{L,\infty|\Phi\Phi}^{(N)}(f, f') &= \frac{1}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \varepsilon_N^{-1+d} \gamma_m(k)^{-1} \\ &\quad \int_{\mathbb{R}^{2d}} d^d x' d^d y' \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}, \\ W_{L,\infty|\text{III}}^{(N)}(f, f') &= \frac{1}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \varepsilon_N^{1+d} \gamma_m(k) \\ &\quad \int_{\mathbb{R}^{2d}} d^d x' d^d y' \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}, \\ W_{L,\infty|\Phi\Pi}^{(N)}(f, f') &= \frac{i}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \varepsilon_N^d \int_{\mathbb{R}^{2d}} d^d x' d^d y' \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f)(x') \varepsilon_N^{-d} (\chi_{[0, \varepsilon_N]^d} * f')(y') e^{ik \cdot (x' - y')}. \end{aligned}$$

Finally, we can take the limit $N \rightarrow \infty$, and we obtain the scaling limit of the two-point functions (after adjusting the scaling dimension),

$$W_{L,\infty|\Phi\Phi}^{(\infty)}(f, f') = \lim_{N \rightarrow \infty} \varepsilon_N^{1-d} W_{m,\infty|\Phi\Phi}^{(N)}(f, f') = \frac{1}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \gamma_m(k)^{-1} \hat{f}(-k) \hat{f}'(k), \quad (5.1)$$

$$W_{L,\infty|\text{III}}^{(\infty)}(f, f') = \lim_{N \rightarrow \infty} \varepsilon_N^{-(1+d)} W_{m,\infty|\text{III}}^{(N)}(f, f') = \frac{1}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \gamma_m(k) \hat{f}(-k) \hat{f}'(k), \quad (5.2)$$

$$W_{L,\infty|\Phi\Pi}^{(\infty)}(f, f') = \lim_{N \rightarrow \infty} \varepsilon_N^{-d} W_{m,\infty|\Phi\Pi}^{(N)}(f, f') = \frac{i}{2(2L)^d} \sum_{k \in \frac{\pi}{L}\mathbb{Z}^d} \hat{f}(-k) \hat{f}'(k), \quad (5.3)$$

as distribution over $C^\infty(\mathbb{T}_L^d)$ (with respect to the usual weak* topology), which are the Wightman two-point functions W_L of the free vacuum state ω_L of continuum scalar field on \mathbb{T}_L^d .

In summary, we find that, although the block-spin scaling map does not define a scaling limit on the algebra $\mathcal{W}_{\infty,L}$, it is still possible to recover the vacuum state of mass m of the continuum scalar field via its Wightman two-point functions W_L defined as scaling limits of the two-point functions $W_{L,M}^{(N)}$ associated with $\omega_{L,M}^{(N)}$ on $\mathcal{W}_{N,L}$. We state this as a theorem:

Theorem 5.1. *Given a sequence of lattice “masses”, $\{\mu_N\}_N$, satisfying (4.18) for some $m > 0$ and the block-spin scaling map, the sequences of two-point functions $\{W_{\mu_{N+M},M}^{(N)}\}_{M \in \mathbb{N}_0}$*

associated with the states $\{\omega_{\mu_{N+M},M}\}_{M \in \mathbb{N}_0}$ on $\mathcal{W}_{N,L}$ for all scales $N \in \mathbb{N}_0$, converge in a scale-coherent way to the free (time-zero) two-point functions of mass m of the continuum scalar field $(\mathcal{W}(\mathfrak{h}_L), \omega_L)$ (see Section 4.2).

Proof. It is well-known that (5.1) – (5.3) are the free (time-zero) two-point functions of mass m over $C^\infty(\mathbb{T}_L^d)$. \square

5.2 Real-space renormalization: Point-like localization

The point-localization renormalization group is a precise formulation of the idea that the algebras $\mathfrak{A}_N(x)$, $x \in \Lambda_N$, correspond to sharply localized operators at $x \in \mathbb{R}^d$ in the continuum. The definition of point-like localized operators is achieved at the level of lattice fields Φ_N and momenta Π_N by the natural inclusion of lattices $\Lambda_N \subset \Lambda_{N'}$ for $N < N'$ as subset of \mathbb{R}^d .

Definition 5.2. The point-localization renormalization group $\{\alpha_{N'}^N\}_{N < N'}$ is the inductive family of *-linear maps defined by the point-localization scaling map between one-particle Hilbert spaces:

$$R_{N+1}^N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{h}_{N+1,L},$$

where

$$R_{N+1}^N(q, p)(x') = 2^d(q, p)(x') \chi_{\Lambda_N \subset \Lambda_{N+1}}(x'), \quad N \in \mathbb{N}_0 \quad (5.4)$$

and

$$R_{N'}^N = R_{N'}^{N'-1} \circ R_{N'-1}^{N'-2} \circ \dots \circ R_{N+1}^N, \quad N < N'.$$

$\Lambda_N \subset \Lambda_{N+1}$ is the canonical inclusion as subsets of \mathbb{R}^d , and $\chi_{\Lambda_N \subset \Lambda_{N+1}}$ is the corresponding characteristic function. The right-hand side of (5.4) is to be understood as an extension of q, p by the zero-function on $\Lambda_{N+1} \setminus \Lambda_N$.

Applying the lattice Fourier transform and its inverse, we find the point-localization scaling map is given by periodic extension in momentum space:

$$\begin{aligned} R_{N+1}^N(\hat{q}, \hat{p})(k') &= \varepsilon_{N+1}^{\frac{d}{2}} \sum_{x' \in \Lambda_{N+1}} 2^d(q, p)(x') \chi_{\Lambda_N \subset \Lambda_{N+1}}(x') e^{-ik' \cdot x'} \\ &= \varepsilon_N^{\frac{d}{2}} 2^{-\frac{d}{2}} \sum_{x \in \Lambda_N} 2^d(q, p)(x) e^{-ik' \cdot x} = 2^{\frac{d}{2}}(\hat{q}, \hat{p})(k'). \end{aligned}$$

In contrast with the previous scaling maps and as a consequence of the rescaling by 2^d in (5.4), we find that $R_{N'}^N$ is only a symplectic homothety for $N < N'$:

$$\sigma_{N'} \circ (R_{N'}^N \times R_{N'}^N) = 2^{d(N'-N)} \sigma_N.$$

Therefore, the induced map $\alpha_{N'}^N : \mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N'}$ is *-compatible linear map but not a *-homomorphism. Nevertheless, we can rescale the symplectic form σ_N determining the Weyl relations to show that for a given $N \in \mathbb{N}_0$ the pullbacks of ground states $\omega_{\mu_{N'}}$ along $\alpha_{N'}^N$ for $N < N'$, which we use to define the scaling limit, yield a sequence of states on $\mathcal{W}_{N,L}$.

Lemma 5.3. *Given a symplectic homothety $R_{N'}^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_{N',L}$ for $N < N'$, the induced *-linear map $\alpha_{N'}^N : \mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N'}$ becomes a *-homomorphism after rescaling the symplectic form according to:*

$$\tilde{\sigma}_{N'} = \lambda_{N,N'}^{-1} \sigma_{N'},$$

where $\lambda_{N,N'} \neq 0$ is the scaling factor of $R_{N'}^N$, i.e. $\sigma_{N'} \circ (R_{N'}^N \times R_{N'}^N) = \lambda_{N,N'} \sigma_N$

Proof. We only need to check the compatibility of $R_{N'}^N$ with the rescaled symplectic form $\tilde{\sigma}_{N'}$ as this implies the compatibility of $\alpha_{N'}^N$ with the Weyl relations (3.1):

$$\tilde{\sigma}_{N'} \circ (R_{N'}^N \times R_{N'}^N) = \lambda_{N,N'}^{-1} \sigma_{N'} \circ (R_{N'}^N \times R_{N'}^N) = \lambda_{N,N'}^{-1} \lambda_{N,N'} \sigma_N = \sigma_N.$$

□

We also recall a general result of Segal about states on Weyl algebras.

Remark 5.4 (Scaled Fock states). *We know from sections 3.1 and 4.1.1, that the symplectic structure σ_N can be expressed as:*

$$\sigma_N((q, p), (q', p')) = -2\Im(\hat{\xi}^{(m)}, \hat{\xi}'^{(m)})_N.$$

Moreover, since $\mathfrak{h}_{N,L} \cong \mathbb{C}^{2r_N}$ is finite dimensional, we know that

$$\|\hat{\xi}^{(m)}\|_N^2 = \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_N} \left(\gamma_m^{(N)}(k)^{-1} |\hat{q}(k)|^2 + \gamma_m^{(N)}(k) |\hat{p}(k)|^2 \right), \quad \xi \in \mathfrak{h}_{N,L},$$

defines non-degenerate (for $m > 0$) norm equivalent to $\|\cdot\|_N$. Thus, by a general theorem of Segal [71], we have a one-parameter family of regular states on the Weyl algebra $\mathcal{W}_{N,L}$:

$$\omega_{m,0}^{(\lambda_N)}(W_N(\xi)) = e^{-\frac{1}{2}\lambda_N \|\hat{\xi}^{(m)}\|_N^2}, \quad \lambda_N \geq 1.$$

The lattice vacuum ω_{μ_N} results from the extremal choice $\lambda_N = 1$ and $\mu_N^2 = \varepsilon_N^2 m^2 + 2d$, cp. (4.3) and (4.4).

From the lemma and the remark we infer the result:

Corollary 5.5. *For $N, M \in \mathbb{N}_0$, the pullback $\omega_{\mu_{N+M},M} = \omega_{\mu_{N+M}} \circ \alpha_{N+M}^N$ defines state on $\mathcal{W}_{N,L}$ explicitly given by:*

$$\omega_{\mu_{N+M}}(\alpha_{N+M}^N(W_N(\xi))) = e^{-\frac{1}{4(2r_N)^d} \sum_{k' \in \Gamma_{N+M}} (\gamma_{\mu_{N+M}}(k')^{-1} |\hat{q}(k')|^2 + \gamma_{\mu_{N+M}}(k') |\hat{p}(k')|^2)},$$

where by a slight abuse of notation \hat{q} and \hat{p} also denote their periodic extensions from Γ_N to Γ_{N+M} .

Proof. Given $N, M \in \mathbb{N}_0$, the lattice vacuum $\omega_{\mu_{N+M}}$ defines a scaled Fock state $\omega_{m,0}^{(\lambda_{N,N+M})}$ on \mathcal{W}_{N+M} with rescaled symplectic form $\tilde{\sigma}_{N+M} = \lambda_{N,N+M}^{-1} \sigma_{N+M}$, where $\lambda_{N,N+M} = 2^{dM}$ and $\mu_{N+M}^2 = \varepsilon_{N+M}^2 m^2 + 2d$ □

Remark 5.6. *In view of the general wavelet scaling map scheme introduced in Section 3.3, the point-localization renormalization group arises from the extremal choice $h_0 = 2^{\frac{d}{2}} \delta_{n,0}$ as low-pass filter. That this choice should correspond to point-like localized field operators in the scaling limit is suggested by $\hat{\phi}(\cdot) = \prod_{N=1}^{\infty} m_0(2^{-N}k) = 1$ which is the inverse Fourier transform of the delta distribution: $\phi_0^{(\varepsilon)} = \varepsilon^{\frac{d}{2}} \delta_0$. Therefore, we expect that the local operators in the continuum are only indirectly accessible through a Wightman-type reconstruction from correlation functions.*

In analogy with the results for the scaling limit in terms of block-spin transformation, the sequences $\{\omega_{\mu_{N+M},M} = \omega_{L,M}^{(N)}\}_{M \in \mathbb{N}_0}$ for $N \in \mathbb{N}_0$ do not converge as we can see from (4.19) and the fact that $\hat{\phi}^{(\varepsilon_N)} = \varepsilon_N^{\frac{d}{2}}$ according to remark 5.6. But, we can use corollary 5.5 to compute the following two-point functions of the time-zero fields Φ_N and momenta Π_N at level N , i.e. for any $x, y \in \Lambda_N$:

$$\begin{aligned}
W_{L,M|\Phi\Phi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{-\frac{1+d}{2}} \Phi_N(x) \varepsilon_N^{-\frac{1+d}{2}} \Phi_N(y))) \\
&= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(t\delta_x^{(N)})) \alpha_{N+M}^N(W_N(s\delta_y^{(N)})))|_{t=s=0} \\
&= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N^{-1} \gamma_{\mu_{N+M}}(k)^{-1} e^{ik \cdot (x-y)}, \\
W_{L,M|\Pi\Pi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{\frac{1-d}{2}} \Pi_N(x) \varepsilon_N^{\frac{1-d}{2}} \Pi_N(y))) \\
&= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(it\delta_x^{(N)})) \alpha_{N+M}^N(W_N(is\delta_y^{(N)})))|_{t=s=0} \\
&= \frac{1}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} \varepsilon_N \gamma_{\mu_{N+M}}(k) e^{ik \cdot (x-y)} \\
W_{L,M|\Phi\Pi}^{(N)}(x, y) &= \omega_{L,M}^{(N)}(\alpha_{N+M}^N(\varepsilon_N^{-\frac{1+d}{2}} \Phi_N(x) \varepsilon_N^{\frac{1-d}{2}} \Pi_N(y))) \\
&= -\partial_t \partial_s \omega_{L,M}^{(N)}(\alpha_{n+M}^N(W_N(t\delta_x^{(N)})) \alpha_{N+M}^N(W_N(is\delta_y^{(N)})))|_{t=s=0} \\
&= \frac{i}{2(2r_N)^d} \sum_{k \in \Gamma_{N+M}} e^{ik \cdot (x-y)},
\end{aligned}$$

Since $\Gamma_N \xrightarrow{N \rightarrow \infty} \frac{\pi}{L} \mathbb{Z}^d$ and comparing with the calculations in Section 5.1, we obtain a result analogous to Theorem 5.1:

Theorem 5.7. *Given a sequence of lattice “masses”, $\{\mu_N\}_{N \in \mathbb{N}_0}$, satisfying (4.18) for some $m > 0$ and the point-localization scaling map, the sequences of two-point functions, $\{W_{\mu_{N+M},M}^{(N)}\}_{M \in \mathbb{N}_0}$, associated with the states, $\{\omega_{\mu_{N+M},M}\}_{M \in \mathbb{N}_0}$ on $\mathcal{W}_{N,L}$ for all scales $N \in \mathbb{N}_0$, converge in a scale-coherent way to the free (time-zero) two-point functions of mass m of the continuum scalar field $(\mathcal{W}(\mathfrak{h}_L), \omega_L)$ (see Section 4.2).*

Proof. The proof works almost identical to that for the block-spin scaling map. The only notable difference is that the convolution with the characteristic function $\varepsilon_N^{-d} \chi_{[0, \varepsilon_N]^d}$ is replaced by a convolution with the delta distribution δ_0 . \square

5.3 Momentum-space renormalization: Sharp momentum cutoff

Since renormalization group schemes in momentum space have a long tradition [81, 33, 82, 87], it is sensible to connect these to the general framework of Section 2.1. While the point-localization scaling map in the previous subsection is induced by the natural inclusion of lattices, $\Lambda_N \subset \Lambda_{N'}$, in real space, the momentum-cutoff renormalization group is obtained from the inclusion of lattices, $\Gamma_N \subset \Gamma_{N'}$, in momentum space. This is made precise in the following definition and precisely corresponds to the idea of taking a partial trace with respect to high-momentum states when coarse graining from states at a fine scale N' to a coarse scale N . Since the momentum cutoff has been used extensively in constructive quantum field theory [39], we also discuss the limit of dynamics in this setting.

Definition 5.8. *The momentum-cutoff renormalization group $\{\alpha_{N'}^N\}_{N < N'}$ is the inductive family of $*$ -homomorphisms defined by the momentum-cutoff scaling map between one-particle Hilbert spaces:*

$$R_{N+1}^N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{h}_{N+1,L},$$

where

$$R_{N+1}^N(\hat{q}, \hat{p})(k') = 2^{\frac{d}{2}}(\hat{q}, \hat{p})(k')\chi_{\Gamma_N \subset \Gamma_{N+1}}(k'), \quad N \in \mathbb{N}_0 \quad (5.5)$$

and

$$R_{N'}^N = R_{N'}^{N'-1} \circ R_{N'-1}^{N'-2} \circ \dots \circ R_{N+1}^N, \quad N < N'.$$

$\Gamma_N \subset \Gamma_{N+1}$ is the canonical inclusion as subsets of $\frac{\pi}{L}\mathbb{Z}^d$, and $\chi_{\Gamma_N \subset \Gamma_{N+1}}$ is the corresponding characteristic function. The right-hand side of (5.5) is to be understood as an extension of \hat{q}, \hat{p} by the zero-function on $\Gamma_{N+1} \setminus \Gamma_N$.

As a trivial consequence of the rescaling by $2^{\frac{d}{2}}$, we have that $R_{N'}^N$ is symplectic for any $N < N'$:

$$\sigma_{N'} \circ (R_{N'}^N \times R_{N'}^N) = \sigma_N,$$

Applying the lattice Fourier transform and its inverse, we find the momentum-cutoff scaling map is given by Fourier interpolation in real space:

$$\begin{aligned} R_{N+1}^N(q, p)(x') &= \varepsilon_{N+1}^{-\frac{d}{2}}(2r_{N+1})^{-d} \sum_{k' \in \Gamma_{N+1}} 2^{\frac{d}{2}}(\hat{q}, \hat{p})(k')\chi_{\Gamma_N \subset \Gamma_{N+1}}(k')e^{ik' \cdot x'} \\ &= \varepsilon_N^{-\frac{d}{2}}(2r_N)^{-d} \sum_{k \in \Gamma_N} (\hat{q}, \hat{p})(k)e^{ik \cdot x'} = \sum_{x \in \Lambda_N} (q, p)(x)(2r_N)^{-d} \sum_{k \in \Gamma_N} e^{-ik \cdot (x-x')}. \end{aligned}$$

The locality of the momentum-cutoff scaling map entails non-locality in real space, which is formally reflected by the use of completely delocalized smearing functions to relate the lattice fields and momenta with their continuum analogues, cp. (3.15) and (3.16):

$$\hat{\Phi}_N(k) = \varepsilon_N^{-\frac{d}{2}}\Phi\left(\sum_{x \in \Lambda_N} \chi_{x+[0, \varepsilon_N]^d} e^{-ik \cdot x}\right), \quad (5.6)$$

$$\hat{\Pi}_N(k) = \varepsilon_N^{-\frac{d}{2}}\Pi\left(\sum_{x \in \Lambda_N} \chi_{x+[0, \varepsilon_N]^d} e^{-ik \cdot x}\right). \quad (5.7)$$

Remark 5.9. *We can translate the momentum-cutoff scaling map into a real-space scaling map by the lattice Fourier transform. This real-space scaling map can be cast into a form analogous to the wavelet scaling map (3.25) with non-local and level-dependent coefficients:*

$$h_n^{(N)} = \left(2^{\frac{1}{2}}r_{N+1}\right)^{-d} \prod_{j=1}^d \frac{\sin\left(\frac{\pi}{2}n_j\right)}{\sin\left(\frac{\pi}{2r_{N+1}}n_j\right)} e^{\frac{i\pi n_j}{2r_{N+1}}}, \quad n \in \{-r_{N+1}, \dots, r_{N+1} - 1\}^d \subset \mathbb{Z}^d.$$

While such a momentum space renormalization group can be used to determine the scaling limit as we show below, the non-locality of the filter makes it ineffective for controlling local operators.

The scaling limit of the free lattice vacua (4.3) can be investigated by the momentum-cutoff renormalization group in a manner similar that for the wavelet method discussed in Section 4.2.2:

$$\omega_{L,M}^{(N)}(W_N(\xi)) = \omega_{\mu_{N+M},M}(W_N(\xi)) = e^{-\frac{1}{4} \left(\|\gamma_{\mu_{N+M}}^{-1/2} \hat{q}_N\|_{\mathfrak{h}_{N,L}}^2 + \|\gamma_{\mu_{N+M}}^{1/2} \hat{p}_N\|_{\mathfrak{h}_{N,L}}^2 \right)}, \quad (5.8)$$

where, for fixed $N \in \mathbb{N}_0$, the dispersion relation $\gamma_{\mu_{N+M}}$ is considered as a function on $\Gamma_N \subset \Gamma_{N+M}$ by restriction. Again, choosing the initial states according to the renormalization condition (4.18) leads to the continuum dispersion relation in the limit $M \rightarrow \infty$. Moreover, the pointwise convergence to continuum dispersion relation implies the weak* convergence of (5.8) to a limit $\omega_{L,\infty}^{(N)}$ which is given by the free vacuum state ω_L of mass m of the continuum scalar field evaluated on Weyl operators with a sharp momentum cut-off at scale $k_{\max} \sim \varepsilon_N^{-1}$:

$$\omega_{L,\infty}^{(N)}(W_N(\xi)) = e^{-\frac{1}{4} \left(\|\gamma_m^{-1/2} \hat{q}_N\|_{\mathfrak{h}_{N,L}}^2 + \|\gamma_m^{1/2} \hat{p}_N\|_{\mathfrak{h}_{N,L}}^2 \right)}. \quad (5.9)$$

Now, it is almost trivial to check that the sequence of limit states, $\{\omega_{L,\infty}^{(N)}\}_{N \in \mathbb{N}_0}$, is projectively consistent for the momentum-cutoff renormalization group,

$$\omega_{L,\infty}^{(N)} = \omega_{L,\infty}^{(N')} \circ \alpha_{N'}^N, \quad N < N',$$

which implies the existence of the scaling limit $\mathcal{W}_{\infty,L}$:

$$\varprojlim_{N \in \mathbb{N}_0} \omega_{L,\infty}^{(N)} = \omega_{L,\infty}^{(\infty)}.$$

In this way, we deduce the following result corresponding to theorems 4.7 and 5.1.

Theorem 5.10. *The GNS representation of $\mathcal{W}_{\infty,L}$ given by the scaling limit $\omega_{L,\infty}^{(\infty)}$ constructed via the momentum-cutoff renormalization group is equivalent to the GNS representation of the continuum scalar field with respect to the free vacuum state ω_L restricted to the subalgebra generated by Weyl operators with finite support in momentum space.*

Proof. By construction the inner product on the GNS Hilbert space $\mathcal{H}_{L,\infty}^{(\infty)}$ is given in terms of the inner product of the inductive limit of one-particle Hilbert spaces $\mathfrak{h}_{\infty,L}$ with dense subspace $\bigcup_{N \in \mathbb{N}_0} \mathfrak{h}_{N,L}$:

$$\omega_{L,\infty}^{(\infty)}(\alpha_{\infty}^N(W_N(\xi))^* \alpha_{\infty}^{N'}(W_{N'}(\xi'))) = e^{-\frac{1}{2}\|\xi\|^2} e^{-\frac{1}{2}\|\xi'\|^2} e^{(\xi,\xi')} = (G(\xi), G(\xi')), \quad (5.10)$$

for $N, N' \in \mathbb{N}_0$ with

$$(\xi, \xi') = \frac{2^{-\frac{d(N+N')}{2}}}{2(2r_0)^d} \sum_{k \in \frac{\pi}{L} \mathbb{Z}^d} \left(\gamma_m(k)^{-1} \overline{\hat{q}_\xi(k)} \hat{q}_{\xi'}(k) + \gamma_m(k) \overline{\hat{p}_\xi(k)} \hat{p}_{\xi'}(k) + i(\overline{\hat{p}_\xi(k)} \hat{q}_{\xi'}(k) - \overline{\hat{q}_\xi(k)} \hat{p}_{\xi'}(k)) \right), \quad (5.11)$$

where we slightly abuse notation to denote by \hat{p}_ξ , \hat{q}_ξ , $\hat{q}_{\xi'}$ and $\hat{p}_{\xi'}$ also their trivial extensions to the infinite lattice $\Gamma_\infty = \frac{\pi}{L} \mathbb{Z}^d$ (cp. (4.5)). On right-hand side of (5.10), $G(\xi) = \exp(\xi)$ is the Glauber vector or coherent state associated with $\xi \in \bigcup_{N \in \mathbb{N}_0} \mathfrak{h}_{N,L}$, and it is known that the map $G : \bigcup_{N \in \mathbb{N}_0} \mathfrak{h}_{N,L} \rightarrow \mathcal{H}_{L,\infty}^{(\infty)}$ is continuous and injective [44, 40]. Now, (5.10)

also shows that the GNS inner product of $\mathcal{H}_{L,\infty}^{(\infty)}$ is identical to that of the symmetric Fock space, $\mathfrak{F}_+(\mathfrak{h}_{\infty,L})$, over $\mathfrak{h}_{\infty,L}$ endowed with (5.11). Because of this we define an embedding of $\iota_L : \mathfrak{h}_{\infty,L} \rightarrow \mathfrak{h}_L$ via the inductive-limit structure of $\mathfrak{h}_{\infty,L}$ analogous to (4.16). Precisely, we define isometries $\iota_L^{(N)} : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_L$ for all $N \in \mathbb{N}_0$:

$$\iota_L^{(N)}(\hat{q}, \hat{p}) = \varepsilon_N^{\frac{d}{2}}(\hat{q}, \hat{p})\chi_{\Gamma_N \subset \Gamma_\infty}.$$

Since we have the identity $\iota_L^{(N')} \circ R_{N'}^N = \iota_L^{(N)}$ for any $N < N'$, we obtain by standard reasoning the sought after ι_L . From the definition of $\iota_L^{(N)}$ it is clear that $\iota_L(\mathfrak{h}_{\infty,L})$ consist of the closure of the subspace of Fourier series with finite support. Therefore, $\iota_L(\mathfrak{h}_{\infty,L}) = \mathfrak{h}_L$ and we have a unitary equivalence of $U_L : \mathfrak{F}_+(\mathfrak{h}_{\infty,L}) \rightarrow \mathfrak{F}_+(\mathfrak{h}_L)$ by functoriality. Similarly, functoriality and the isometricity of $\iota_L^{(N)}$ for each $N \in \mathbb{N}_0$, implies the existence of an injective *-homomorphism $\Phi_L : \mathcal{W}_{\infty,L} \rightarrow \mathcal{W}_L$ such that

$$\pi_{L,\infty}^{(\infty)}(W) = U_L^* \pi_{\omega_L}(\Phi_L(W)) U_L, \quad \forall W \in \mathcal{W}_{\infty,L}.$$

□

To conclude our discussion about the momentum-cutoff scaling map, we also take a look at the limit of dynamics. We observe that this limit is considerably simpler with respect to this scheme compared to that in the wavelet scheme (see Section 4.2.4), i.e. the limit of dynamics exist even in the sense of Hamiltonians as depicted in figure 1.

At the level of the one-particle space $\mathfrak{h}_{N,L} \cong \ell^2(\Gamma_N, (2r_N)^{-d} \mu_{\Gamma_N})$, the action of $H_{L,0}^{(N)}$ is (essentially) given by multiplication with the dispersion relation γ_{μ_N} (cp. remark 4.10). Since the momentum-cutoff scaling map is (up to rescaling) induced by the inclusion of lattices $\Gamma_N \subset \Gamma_{N'}$ for $N < N'$, the action of $H_{L,0}^{(N')}$ restricts consistently to the image of $\mathcal{H}_{N,L}$ inside $\mathcal{H}_{N',L}$ for $N < N'$ because the aforesaid multiplication preserves the support in momentum space. This observation leads to the following result.

Theorem 5.11. *Given the momentum-cutoff renormalization group $\{\alpha_{N'}^N\}_{N < N'}$ and a sequence of lattice “masses” $\{\mu_N\}_{N \in \mathbb{N}_0}$ satisfying condition (4.18) with a physical mass $m > 0$, we obtain a sequence of (self-adjoint) renormalized Hamiltonians $\{H_{L,M}^{(N)}\}_{M \in \mathbb{N}_0}$ on $\mathcal{H}_{N,L}$ for each $N \in \mathbb{N}_0$. Moreover, each sequence converges in the norm resolvent sense,*

$$H_{L,M}^{(N)} \xrightarrow{\text{nR}} H_{L,\infty}^{(N)},$$

to a self-adjoint operator $H_{L,\infty}^{(N)}$ and the sequence $\{H_{L,\infty}^{(N)}\}_{N \in \mathbb{N}_0}$ defines a self-adjoint inductive limit,

$$\varinjlim_{N \in \mathbb{N}_0} H_{L,\infty}^{(N)} = H_{L,\infty}^{(\infty)},$$

on the GNS Hilbert space $\mathcal{H}_{L,\infty}^{(\infty)}$. The Hamiltonian $H_{L,\infty}^{(\infty)}$ of the scaling limit $\omega_{L,\infty}^{(\infty)}$ is identified with the free Hamiltonian H_L of mass m on \mathcal{H}_L by theorem 5.10.

Proof. We realize the GNS Hilbert space $\mathcal{H}_{L,\infty}^{(\infty)}$ as an inductive limit of a sequence of the Hilbert spaces $\{\mathcal{H}_{N,L}^{(\infty)}\}_{N \in \mathbb{N}_0}$ by applying the GNS construction with respect to the restriction $\omega_{L,\infty}^{(\infty)}$ to each $\mathcal{W}_{N,L}$. By construction, the inductive system of isometries $V_{N'}^N : \mathcal{H}_{N,L}^{(\infty)} \rightarrow \mathcal{H}_{N',L}^{(\infty)}$, $N < N'$, is induced by the momentum-cutoff renormalization group. Applying

the GNS construction also to the renormalized states (5.8), we obtain another sequence of Hilbert spaces, $\{\mathcal{H}_{N,L}^{(M)}\}_{M \in \mathbb{N}_0}$, for each $N \in \mathbb{N}_0$. By von Neumann's uniqueness theorem the representations of $\mathcal{W}_{N,L}$ on all $\mathcal{H}_{N,L}^{(M)}$ and $\mathcal{H}_{N,L}^{(\infty)}$ are unitarily equivalent to the representation on $\mathcal{H}_{N,L}$, because all the GNS Hilbert spaces are symmetric Fock space over the finite-dimensional vector space $\ell^2(\Gamma_N, (2r_N)^{-d} \mu_{\Gamma_N})$ with inner products given in terms of either some $\gamma_{\mu_{N+M}}$ or γ_m which are bounded above and strictly bounded away from zero on Γ_N ¹⁴. Therefore, we implicitly identify all the GNS Hilbert spaces with $\mathcal{H}_{N,L}$ and define the renormalized Hamiltonians $H_{L,M}^{(N)}$ by restriction of $H_{L,0}^{(N+M)}$ to $\mathcal{H}_{N,L}^{(M)}$ inside $\mathcal{H}_{N+M,L}$, i.e.:

$$H_{L,M}^{(N)} = H_{L,0|\mathcal{H}_{N,L} \subset \mathcal{H}_{N+M,L}}^{(N+M)}.$$

Each $H_{L,M}^{(N)}$ is self-adjoint as it arises as the second quantization of a self-adjoint multiplication operator (with $\gamma_{\mu_{N+M}}$) on $\mathfrak{h}_{N,L}$. Analogously, we define $H_{L,\infty}^{(N)}$ as the self-adjoint second quantization of the self-adjoint multiplication operator by γ_m on $\mathfrak{h}_{N,L}$. The uniform convergence $\gamma_{\mu_{N+M}} \rightarrow \gamma_m$ on Γ_N implies the convergence $H_{L,M}^{(N)} \rightarrow H_{L,\infty}^{(N)}$ in the norm resolvent sense, cp. [49]. Using the identification of $\mathcal{H}_{N,L}$ with $\mathcal{H}_{N,L}^{(\infty)}$, we observe that the sequence $\{H_{L,\infty}^{(N)}\}_{N \in \mathbb{N}_0}$ satisfies the compatibility condition:

$$H_{L,\infty}^{(N')} V_{N'}^N = V_{N'}^N H_{L,\infty}^{(N)}, \quad \forall N < N'.$$

Moreover, the dense subspaces $\mathcal{D}_{N,L}^{(\infty)} \subset \mathcal{H}_{N,L}^{(\infty)}$, $N \in \mathbb{N}_0$, of vectors with finite particle number are cores for each $H_{L,\infty}^{(N)}$ and satisfy $V_{N'}^N \mathcal{D}_{N,L}^{(\infty)} \subset \mathcal{D}_{N',L}^{(\infty)}$. Therefore, the inductive-limit operator $H_{L,\infty}^{(\infty)}$ exists and is essentially self-adjoint on the algebraic inductive limit of the subspaces $\mathcal{D}_{N,L}^{(\infty)}$ which is dense in $\mathcal{H}_{L,\infty}^{(\infty)}$. It is easy to check that this operators agrees with the free Hamiltonian H_L of mass m on \mathcal{H}_L . \square

6 Consequences and Outlook

To conclude the article, we discuss further implications of the results presented in Sections 2 – 5 and potential lines of future research. So far, we have focused on scaling limits of free lattice ground states of scalar quantum fields as these form a principal family of basic models on which to build more complicated ones, and therefore are required to be under best possible control. But, the general renormalization group scheme presented in Section 2 as well as the more specific wavelet method are not limited to such a setting, thus leading among other things to the following rough division into topics allowing for generalizations and extensions:

- interactions (ground states and dynamics) and connections with Euclidean methods,
- fermions and connections with conformal field theory,
- scaling limits of thermal (KMS) states and applications to quantum statistical mechanics,

¹⁴Alternatively, it also easy to check that the Bogoliubov transformations which realize the equivalence are unitarily implemented.

- wavelets with non-linear domains (curved spacetimes, e.g. deSitter) and/or target spaces (gauge field, sigma models).

Below we discuss in more detail some of the raised topics.

6.1 Other continuum models and locality from lattices

First of all, we considered only massive free fields in this paper. One difference that appears with massless fields is the zero mode: the inverse of the dispersion relation is not defined at $k = 0$, and this must be avoided somehow (possibly by excluding the zero-mode before taking the infinite volume limit or by enforcing anti-periodic (twisted) boundary conditions). In addition, massless free fields with different helicity might require different symplectic structures [55].

In view of the classical construction of interacting scalar fields in 1+1-dimensional spacetimes [39], we point out that the general concept of limit dynamics (2.13) together with the momentum-cutoff renormalization group (see Section 5.3) can be used to recover the existence of interacting dynamics on the continuum time-zero net of the scalar field with respect to free vacuum. Clearly, this observation should be generalized in two aspects: First, it should be possible to recover the interacting dynamics using the wavelet renormalization group as well, which would be advantageous from the point of view of locality. Second, it would be conceptually preferred (and necessary for $d > 1$) to work with a scaling limit of the interacting lattice ground states instead of the free lattice ground states as the use of the latter is only possible because of the local Fock property in two-dimensional spacetime [37].

If we consider more general models than lattice free fields, we hope to obtain (interacting) quantum field theories by the operator-algebraic renormalization procedure explained in Section 2.1. A general justification of this expectation is due to the result on finite propagation speed of the scaling limit dynamics via Lieb-Robinson bounds for the approximate dynamics on the lattice, cp. [65, 13]. The main reason why a finite propagation speed can be deduced from these bounds in the scaling limit of the lattice free field is due to the correct scaling of the Lieb-Robinson velocity (with respect to the lattice spacing ε) and a subexponential bound on the field-strength renormalization, i.e. the rescaling of the one-particle vectors q, p with respect to the lattice spacing ε . Therefore, if said scaling is preserved in the presence of interaction on the lattice, a finite propagation speed in the scaling limit will be attained. Explicit Lieb-Robinson bounds for discretized scalar fields possibly with interaction are available in [62], see also [61].

Thus, as also suggested in [13], this might provide a way to approximate local spacetime algebras at the lattice level by adapting the definition of these algebras in the Hamiltonian setting (4.28) to the discretized setting, cp. [38, Chapter IV, Definition 2.8].:

$$\mathcal{A}_{N,L}(\mathcal{O}) = \left(\bigcup_{t \in \mathbb{R}} \eta_{L|t}^{(N)} (\mathcal{W}_{N,L}(\mathcal{O}(t) \cap \Lambda_N)) \right)'' , \quad N \in \mathbb{N}_0.$$

$\mathcal{O} \subset \mathbb{R} \times \mathbb{T}_L^d$ should be a suitable open bounded subset with time slices $\mathcal{O}(t)$, $t \in \mathbb{R}$. $\mathcal{W}_{N,L}(\mathcal{O}(t) \cap \Lambda_N) \subset \mathcal{W}_{N,L}$ is the Weyl subalgebra of the sublattice $\mathcal{O}(t) \cap \Lambda_N$ (extension by $\mathbb{1}$ outside). $\eta_L^{(N)} : \mathbb{R} \curvearrowright \mathcal{W}_{N,L}$ is the approximate dynamics at level $N \in \mathbb{N}_0$.

6.2 Extension to fermions

Similar to the bosonic case, the purely kinematical part of the renormalization group scheme of Section 2.1 can be treated by first and second quantization for fermions, cp. [30, 42, 84] for a related construction in the context of MERA. Specifically, if we use dyadically scaled family $\{\Lambda_N\}_{N \in \mathbb{N}_0}$ of a basic lattice $\Lambda_0 = \varepsilon_0 \{-r_0, \dots, 0, \dots -r_0 - 1\}^d$ serving as discretizations of the torus \mathbb{T}_L^d as before, we can introduce at each scale N the one-particle spaces,

$$\mathfrak{h}_{N,L} = \ell^2(\Lambda_N) \otimes \mathbb{C}^s,$$

with s denoting the number of spinor components (e.g. $s = 2$ for $d = 1$, $s = 4$ for $d = 3$ for Dirac fermions). Then fermionic second quantization provides the quantum systems

$$\mathfrak{A}_N = \mathfrak{A}_{\text{CAR}}(\mathfrak{h}_{N,L}), \quad \mathcal{H}_{N,L} = \mathfrak{F}_-(\mathfrak{h}_{N,L}),$$

where $\mathfrak{A}_{\text{CAR}}(\mathfrak{h})$ denotes the C^* -algebra with one-particle Hilbert space \mathfrak{h} based on the complex form of the canonical anti-commutation relations [26], and $\mathfrak{F}_-(\mathfrak{h})$ is the associated anti-symmetric Fock space. Since the algebras \mathfrak{A}_N are generated by an anti-linear map (the dimensionless, complex fermion field),

$$\psi_N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{A}_{\text{CAR}}(\mathfrak{h}_{N,L}),$$

$$\{\psi_N(\xi), \psi_N(\eta)\} = 0, \quad \{\psi_N(\xi), \psi_N^\dagger(\eta)\} = (\xi, \eta)_{\mathfrak{h}_{N,L}} \mathbb{1},$$

with $\xi, \eta \in \mathfrak{h}_{N,L}$ and $\psi_N^\dagger(\eta) = \psi_N(\eta)^*$, it is clear that the wavelet renormalization group can be adapted at the one-particle level by (cf. Section 3.3):

$$R_{N+1}^N : \mathfrak{h}_{N,L} \longrightarrow \mathfrak{h}_{N+1,L}, \quad R_{N+1}^N(\xi) = \sum_{x \in \Lambda_N} \xi(x) \sum_{n \in \mathbb{Z}^d} h_n \delta_{x+\varepsilon_N n}^{(N+1)},$$

which yields an inductive system of Hilbert spaces,

$$\varinjlim_{N \in \mathbb{N}_0} \mathfrak{h}_{N,L} = \mathfrak{h}_{\infty,L} = L^2(\mathbb{T}_L^d), \quad (6.1)$$

because $R_{N+2}^{N+1} \circ R_{N+1}^N = R_{N+2}^N$ and $(R_{N+1}^N(\xi), R_{N+1}^N(\eta))_{\mathfrak{h}_{N+1,L}} = (\xi, \eta)_{\mathfrak{h}_{N,L}}$ (for orthogonal wavelets). Then, we relate the fermion fields at different scales,

$$\alpha_{N+1}^N(\psi_N(\xi)) = \psi_{N+1}(R_{N+1}^N(\xi)),$$

thereby defining an inductive system of C^* -algebras:

$$\varinjlim_{N \in \mathbb{N}_0} \mathfrak{A}_N = \mathfrak{A}_\infty = \mathfrak{A}_{\text{CAR}}(L^2(\mathbb{T}_L^d)). \quad (6.2)$$

We note that the second equalities in (6.1) and (6.2) are a major departure from the bosonic case which are essentially due to the fact that the one-particle spaces carry a unitary structure instead of a symplectic structure and the continuity of the CAR functor, i.e. $\mathfrak{A}_{\text{CAR}}(\mathfrak{k}) = \mathfrak{A}_{\text{CAR}}(\mathfrak{h})$ for $\mathfrak{k} = \mathfrak{h}$. Moreover, we observe that the asymptotic embeddings $R_\infty^N : \mathfrak{h}_{N,L} \rightarrow \mathfrak{h}_{\infty,L}$ and $\alpha_\infty^N : \mathfrak{A}_N \rightarrow \mathfrak{A}_\infty$ are explicitly realized by:

$$R_\infty^N(\xi) = \xi *_{\Lambda_N} \phi^{(\varepsilon_N)}, \quad \alpha_\infty^N(\psi_N(\xi)) = \psi(\xi *_{\Lambda_N} \phi^{(\varepsilon_N)}),$$

for a choice of orthogonal scaling function $\phi \in L^2(\mathbb{T}_L^d)$ and the continuum fermion field ψ .

We close our preliminary treatment of fermions with an observation concerning the construction of scaling limits of initial families of states, $\omega_{L,0}^{(N)}$, in the quasi-free setting (covering the case of free ground states for lattice fermions): It is known [26], that a quasi-free state on $\mathfrak{A}_{\text{CAR}}(\mathfrak{h})$ is determined by its two-point,

$$\omega(\psi(\xi)\psi^\dagger(\eta)) = (\xi, T(\eta))_{\mathfrak{h}}, \quad \xi, \eta \in \mathfrak{h}, \quad (6.3)$$

for some operator, $T : \mathfrak{h} \rightarrow \mathfrak{h}$, with $0 \leq T \leq 1$. Thus, if we consider a family of states $\{\omega_{L,0}^{(N)}\}_{N \in \mathbb{N}_0}$ parametrized by a family of operators $\{T_L^{(N)}\}_{N \in \mathbb{N}_0}$, the existence of a scaling limit $\omega_{L,\infty}^{(\infty)}$ can be rephrased as the weak convergence:

$$\text{w-lim}_{M \rightarrow \infty} T_{L,M}^{(N)} = T_{L,\infty}^{(N)}, \quad \forall N \in \mathbb{N}_0, \quad (6.4)$$

with $T_{L,M}^{(N)} = (R_{N+M}^N)^* T_L^{(N+M)} R_{N+M}^N$, which will define an operator $T_{L,\infty}^{(\infty)}$ on $L^2(\mathbb{T}_L^d)$ satisfying $0 \leq T_{L,\infty}^{(\infty)} \leq 1$ by Proposition 2.2. Thus, as long as the Fourier transform of the kernel of the operator $T_L^{(N)}$ has appropriately bounded growth uniformly in N , we can adopt Lemma 4.2 to conclude convergence of (6.4).

6.3 Connections with multi-scale entanglement renormalization (MERA)

We note that there is a natural decomposition of the wavelet scaling map (3.25) into a tensor-product embedding of the form $W_N(\xi) \mapsto W_N(\xi) \otimes \mathbf{1}_{\mathcal{H}_{N+1|N}}$ and the adjoint action by a unitary $U_{N+1} \in \mathcal{U}(\mathcal{H}_{N+1,L})$, i.e.:

$$\alpha_{N+1}^N(W_N(\xi)) = \text{Ad}_{U_{N+1}}(W_N(\xi) \otimes \mathbf{1}_{\mathcal{H}_{N+1|N}}). \quad (6.5)$$

Here, we denote by $\mathcal{H}_{N+1|N} = \mathfrak{F}_+(\ell^2(\Lambda_{N+1} \setminus \Lambda_N))$ the natural tensor-product complement of $\mathcal{H}_{N,L}$ inside $\mathcal{H}_{N+1,L}$. Thus, the wavelet scaling map belongs to the class of examples alluded to in (2.7). This follows from the decomposition of R_{N+1}^N into a symplectic automorphism of $\mathfrak{h}_{N+1,L}$ and the embedding $\ell^2(\Lambda_N) \rightarrow \ell^2(\Lambda_{N+1})$ defined by the inclusion $\Lambda_N \subset \Lambda_{N+1}$ as well as (multiplicative) second quantization:

$$\alpha_{N+1}^N = \mathfrak{F}_+(R_{N+1}^N) = \mathfrak{F}_+(S_{N+1}) \circ \mathfrak{F}_+(\ell^2(\Lambda_N) \rightarrow \ell^2(\Lambda_{N+1})) = \text{Ad}_{U_{N+1}} \circ (\mathcal{W}_{N,L} \rightarrow \mathcal{W}_{N+1,L}).$$

The symplectic automorphism S_{N+1} is determined by the scaling equation (3.39),

$$S_{N+1}(x', y') = \sum_{n \in \mathbb{Z}^d} h_n \delta_{\epsilon_{N+1}n, x' - y'}, \quad (6.6)$$

as a kernel relative to the standard basis of $\ell^2(\Lambda_{N+1})$.

From this we can make three basic observations, see also [14, 13]:

- The C^* -inductive limit $\mathcal{W}_{\infty,L}$ is $*$ -isomorphic to an infinite (minimal) tensor-product of the basic algebra $\mathcal{W}(L^2(\mathbb{R}))$. This follows from an inductive construction using the unitary conjugacy of the renormalization group $\{\alpha_{N'}^N\}_{N' > N}$ with the tensor-product embedding, cp. (6.5).
- A projectively-consistent family of states $\{\omega_{L,\infty}^{(N)}\}_{N \in \mathbb{N}_0}$ or alternatively a state $\omega_{L,\infty}^{(\infty)}$ as in the definition of the scaling limit (2.11) on $\mathcal{W}_{\infty,L}$ is analogous to a MERA (see

e.g. [75, 29]), i.e. the GNS vectors of $\{\Omega_N\}_{N \in \mathbb{N}_0}$ given by the projective family or the corresponding restrictions of $\omega_{L,\infty}^{(\infty)}$ satisfy, see Section 2.2:

$$V_{N'}^N \Omega_N = \Omega_{N'}, \quad N < N', \quad (6.7)$$

where $V_{N'}^N$ is the GNS (partial) isometry induced by $\alpha_{N'}^N$. The connection of the latter with a MERA follows from the observation that a partial isometry compatible with (6.5) is given by:

$$\psi_N \mapsto U_{N+1}(\psi_N \otimes \Omega_{N+1|N}^{(0)}), \quad (6.8)$$

leading to a representation of $\mathcal{W}_{\infty,L}$. Here, $\Omega_{N+1|N}^{(0)}$ is the Fock vacuum of the space $\mathfrak{F}_+(\ell^2(\Lambda_{N+1} \setminus \Lambda_N))$, and the MERA is given by the partial isometry $I_{N+1}^N : \mathfrak{F}_+(\ell^2(\Lambda_N)) \rightarrow \mathfrak{F}_+(\ell^2(\Lambda_{N+1}))$, $I_{N+1}^N(\psi_N) = \psi_N \otimes \Omega_{N+1|N}^{(0)}$, and the unitary disentangler U_{N+1} . In this sense (6.7) generalizes (6.8) subsuming U_{N+1} and I_{N+1}^N by the partial isometry $V_{N'}^N$ coming from the GNS construction.

- Each α_{N+1}^N extends to a normal, unital *-morphism from $B(\mathcal{H}_{N,L})$ to $B(\mathcal{H}_{N+1,L})$, which allows us to consider the inductive limit $\varinjlim_{N \in \mathbb{N}_0} B(\mathcal{H}_{N,L}) = B_\infty$. Thus, as long as we are interested in properties pertaining to the scaling limit $\omega_{L,\infty}^{(\infty)}$ and the weak closure of the associated GNS representation of a family of regular initial states $\omega_{L,0}^{(N)}$, we can substitute B_∞ for \mathcal{W}_∞ in the analysis.

6.4 Scaling maps and biorthogonal wavelets

In the definition of the wavelet scaling map (see Definition 3.3) there is no a priori need to choose the same low-pass filter $\{h_n\}_{n \in \mathbb{Z}^d}$ for the complementary real, Lagrangian subspaces referred to by the decomposition $\xi = \varepsilon_N^{\frac{d+1}{2}} q + i\varepsilon_N^{\frac{d-1}{2}} p$ for $\xi \in \mathfrak{h}_{N,L}$. More precisely, the scaling map $R_{N'}^N$ will be symplectic as long as the biorthogonality condition [23],

$$\sum_{n \in \mathbb{Z}^d} h_n^{(q)} h_{n+2k}^{(p)} = \delta_{k,0}, \quad k \in \mathbb{Z}^d, \quad (6.9)$$

is satisfied for two low-pass filters $\{h_n^{(q)}\}_{n \in \mathbb{Z}^d}$ and $\{h_n^{(p)}\}_{n \in \mathbb{Z}^d}$, each associated the component q and p separately, cp. Lemma 3.4. It is evident that the results proved for the wavelet scaling map remain true in the biorthogonal setting, with the advantage that the decay properties of scaling functions can be independently adjusted to the time-zero fields and momenta. Moreover, the biorthogonal setting allows for a greater freedom regarding symmetry properties of the scaling functions $\phi, \check{\phi}$ if compact support is required for the latter [23, Ch. 8]. More generally, we could also relax the condition that the low-pass filter $\{h_n\}_{n \in \mathbb{Z}^d}$ is finite, thereby also allowing for the possibility of smooth wavelets [58] albeit at the cost of loosing the strict spatial locality structure in the inductive limit.

6.5 Entropy and Type I approximation of local algebras

In a quantum theory with finitely many degrees of freedom physically relevant states ω are defined by density matrices ρ_ω on a Hilbert space \mathcal{H} as $\omega = \text{Tr}(\rho_\omega \cdot)$, and the von Neumann entropy is given by $S_{\text{vN}}(\omega) = -\text{Tr}(\rho_\omega \ln \rho_\omega)$. It is well known that this definition has the

problem to be infinite on the local algebras appearing in QFT. Yet, the continuum local algebras can be approximated by lattice algebras by Theorem 4.7 and Theorem 4.22 and lattice algebras are manifestly type I by the Stone-von Neumann uniqueness theorem for the CCR algebras with finitely many degrees of freedom. In principle, this allows for computing the von Neumann entropy of lattice algebras with respect to the Minkowski vacuum state and to study how it diverges as one approaches the continuum algebra (cf. the use of intermediate type I factors [54]).

The question of computing the von Neumann entropy is also connected with modular theory and, thereby, the recovery of symmetries in the continuum and infinite volume limits. For instance, recall that in Minkowski space, the modular group of the algebra $\mathcal{A}(W_1)$ of the Rindler wedge W_1 with respect to the vacuum vector Ω is particularly important since, when the Bisognano-Wichmann property holds, it has a geometric meaning given by

$$U(\Lambda_{W_1}(2\pi t)) = \Delta_{\mathcal{A}(W_1), \Omega}^{-it}, \quad (6.10)$$

where Λ_{W_1} is the one-parameter group of boosts in the x_1 -direction (preserving W_1) and U is the covariant unitary positive energy representation of the Poincaré group. But, the modular operator $\Delta_{\mathcal{A}(W_1), \Omega}$ also formally encapsulates the density matrix of the thermal state $\omega|_{\mathcal{A}(W_1)}$. It is well known that (6.10) holds in any theory generated by Wightman fields [6] and it can also be derived from more general structural assumptions [60, 25]. On the other hand, for a fixed state and a growing sequence of von Neumann algebras, the modular groups of the union can be approximated by the modular group of the subalgebras [53, Lemma 3]. This raises an interesting question of what the modular groups of the lattice algebras are, and how the Bisognano-Wichmann property in the continuum is restored. This is also connected with the question of computing entropy of local algebras on lattices, as the density matrix is obtained by tracing out the modular operator to the local algebra

6.6 UV vs. IR scaling limits: Momentum transfer to small scales

The momentum space setting of Section 5.3 is suitable to further discuss distinctive aspects between ultraviolet and infrared scaling limits raised in Section 1. To be rather specific, we contrast two operator-algebraic schemes: the scaling algebra approach by Buchholz and Verch [15, 16], which aims at the construction of ultraviolet scaling limits of continuum field theories, and our setting that primarily deals with the construction of continuum models via infrared scaling limits of lattice models. The differences becomes manifest at a technical level by modifying the momentum-cutoff renormalization group of Definition 5.8 to arrive at another renormalization group that resembles the scaling map introduced in [16, Eq. (2.7)] for continuum scalar fields. This modification is given in terms of a shift of support of one-particle vectors $\hat{q}, \hat{p} \in \mathfrak{h}_{N,L}$ towards higher momentum (apart from the zero mode $k = 0$) via the canonical inclusion of dual lattices $2\Gamma_N \subset \Gamma_{N'}$ for $N < N'$,

$$R_{N+1}^N(\hat{q}, \hat{p})(k') = (2^{\frac{d+1}{2}}\hat{q}, 2^{\frac{d-1}{2}}\hat{p})(2^{-1}k')\chi_{2\Gamma_N \subset \Gamma_{N+1}}(k'), \quad N \in \mathbb{N}_0 \quad (6.11)$$

thereby implementing a ultraviolet scaling limit procedure in the general setting of Section 4.

By applying the lattice Fourier transform, we see that the real-space version is the following, where it is evident that these maps are symplectic:

$$R_{N+1}^N(q, p)(x') = (2^{\frac{1}{2}}q, 2^{-\frac{1}{2}}p)(2x').$$

The latter can be directly compared with the scaling map (up to an additional factor of $2^{\frac{d}{2}}$ due to periodic boundary conditions on the lattice),

$$R_{N+1}^N(f)(x') = 2^{\frac{d+1}{2}} \Re(f)(2x') + i2^{\frac{d-1}{2}} \Im(f)(2x'),$$

used in [16] for complex one-particle vectors $f \in \mathcal{S}(\mathbb{R}^d)$ in the continuum subject to the symplectic form (4.14). By applying the resulting renormalization group to the lattice vacua $\{\omega_{\mu_N}^{(N)}\}_{N \in \mathbb{N}_0}$, we find the following transformation behavior:

$$\omega_{\mu_{N+M}, M}^{(N)}(W_N(\xi)) = \omega_{\mu_{N+M}}^{(N+M)}(\alpha_{N+M}^N(W_N(\xi))) = \omega_{\mu_{N+M}}^{(N)}(W_N(\xi)).$$

Here, we explicitly indicate the scale $N \in \mathbb{N}_0$ to which the state is associated by the upper index (N) . In other words, the action of the momentum-transfer renormalization group identifies the renormalized state $\omega_{L, M}^{(N)} = \omega_{\mu_{N+M}, M}^{(N)}$ at scale N after M steps with the lattice vacuum $\omega_{\mu_{N+M}}^{(N)}$ of lattice mass μ_{N+M} at scale N . Thus, assigning a physical mass $m > 0$ to the lattice mass μ_N uniformly among all scales N according to (4.4), this entails (at least for $d > 1$):

$$\omega_{L, M}^{(N)}(W_N(\xi)) = \omega_{2^{-M}m, 0}^{(N)}(W_N(\xi)). \quad (6.12)$$

In this respect, the relation between lattice mass μ_N and physical mass m , the momentum-transfer renormalization group universally¹⁵ leads to the family of massless lattice vacua $\{\omega_{0,0}^{(N)}\}_{N \in \mathbb{N}_0}$ as the projective family defining the scaling limit, i.e. (4.4) in the limit $m \rightarrow 0+$. At first sight, this appears to be similar to the observation made in [16] for ultraviolet scaling limits of massive continuum free fields (in $d > 1$), and it is tempting to assume that the scaling limit of (6.12) equals the vacuum of the massless continuum free field in finite volume ω_L . But, this is not as evident as for the wavelet renormalization group because the relation between the scaling-limit algebra $\mathcal{W}_{\infty, L}$ and the continuum algebra \mathcal{W}_L remains rather obscure.

In summary, we conclude that while there are technical similarities between the scaling algebra approach of [15, 16], the latter should be distinguished from the approach detailed here. Moreover, although it is possible to use the general method of Section 2.1 to implement ultraviolet scaling limits as explicated above, it is conceptually unlikely to be meaningful in the context of scaling limits of lattice models as in Section 4, where the goal is to obtain an infrared scaling limit.

References

- [1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] H. Araki. von Neumann algebras of local observables for free scalar field. *J. Mathematical Phys.*, 5:1–13, 1964. ISSN 0022-2488. <https://dx.doi.org/10.1063/1.1704063>.
- [3] H. Araki and S. Yamagami. On quasi-equivalence of quasifree states of the canonical commutation relations. *Publ. Res. Inst. Math. Sci.*, 18(2):703–758 (283–338), 1982. ISSN 0034-5Di318. doi: 10.2977/prims/1195183576. URL <https://doi.org/10.2977/prims/1195183576>.

¹⁵This will hold more generally as long as $\lim_{M \rightarrow \infty} 2^{-2M} \varepsilon_{N+M}^{-2} (\mu_{N+M}^2 - 2d) = 0$ for all $N \in \mathbb{N}_0$.

- [4] G. Battle. *Wavelets and Renormalization*, volume 10 of *Series in Approximations and Decompositions*. World World Scientific Publishing Company, 1999. doi: 10.1142/3066. URL <https://doi.org/10.1142/3066>.
- [5] C. Bény and T. J. Osborne. Information-geometric approach to the renormalization group. *Physical Review A - Atomic, Molecular, and Optical Physics*, 92(2):022330, 2015. doi: 10.1103/PhysRevA.92.022330. URL <https://doi.org/10.1103/PhysRevA.92.022330>.
- [6] J. J. Bisognano and E. H. Wichmann. On the duality condition for quantum fields. *J. Mathematical Phys.*, 17(3):303–321, 1976. ISSN 0022-2488. doi: 10.1063/1.522898. URL <https://doi.org/10.1063/1.522898>.
- [7] B. E. Blackadar. *Operator Algebras: Theory of C*-Algebras and von Neumann Algebras*, volume 122 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag Berlin Heidelberg, 2006. ISBN 978-3-540-28486-4. doi: 10.1007/3-540-28517-2. URL <https://doi.org/10.1007/3-540-28517-2>.
- [8] C. Borgs. Confinement, deconfinement and freezing in lattice Yang-Mills theories with continuous time. *Communications in Mathematical Physics*, 116(2):309–342, 1988. ISSN 0010-3616. doi: 10.1007/bf01225259. URL <https://doi.org/10.1007/bf01225259>.
- [9] H. Bostelmann, C. D’Antoni, and G. Morsella. Scaling algebras and point-like fields: A nonperturbative approach to renormalization. *Communications in Mathematical Physics*, 285:763–798, 2009. doi: 10.1007/s00220-008-0613-3. URL <https://doi.org/10.1007/s00220-008-0613-3>.
- [10] O. Bratteli and D. W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 1: C*-and W*-Algebras, Symmetry Groups, Decomposition of States*. Theoretical and Mathematical Physics. Springer, second edition, 1987. doi: 10.1007/978-3-662-02520-8. URL <https://doi.org/10.1007/978-3-662-02520-8>.
- [11] O. Bratteli and D. W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 2: Equilibrium States, Models in Quantum Statistical Mechanics*. Theoretical and Mathematical Physics. Springer, second edition, 1997. doi: 10.1007/978-3-662-03444-6. URL <https://doi.org/10.1007/978-3-662-03444-6>.
- [12] G. K. Brennen, P. Rohde, B. C. Sanders, and S. Singh. Multiscale quantum simulation of quantum field theory using wavelets. *Physical Review A: Atomic, Molecular, and Optical Physics*, 92(3):032315, 2015. ISSN 1050-2947. doi: 10.1103/physreva.92.032315. URL <https://doi.org/10.1103/physreva.92.032315>.
- [13] A. Brothier and A. Stottmeister. Canonical Quantization of 1+1-dimensional Yang-Mills Theory: An Operator Algebraic Approach. *Preprint, arXiv: 1907.05549*, 2019. URL <http://arxiv.org/abs/1907.05549>.
- [14] A. Brothier and A. Stottmeister. Operator-algebraic Construction of Gauge Theories and Jones’ Actions of Thompson’s Groups. *Communications in Mathematical Physics*, 2019. doi: 10.1007/s00220-019-03603-4. URL <https://doi.org/10.1007/s00220-019-03603-4>.

- [15] D. Buchholz and R. Verch. Scaling Algebras and Renormalization Group in Algebraic Quantum Field Theory. *Reviews in Mathematical Physics*, 7(8): 1195–1239, 1995. ISSN 0129-055X. doi: 10.1142/s0129055x9500044x. URL <https://doi.org/10.1142/S0129055X9500044X>.
- [16] D. Buchholz and R. Verch. Scaling Algebras and Renormalization Group in Algebraic Quantum Field Theory. *Reviews in Mathematical Physics*, 10(6): 775–800, 1998. ISSN 0129-055X. doi: 10.1142/s0129055x98000252. URL <https://doi.org/10.1142/S0129055X98000252>.
- [17] D. Buchholz and E. H. Wichmann. Causal independence and the energy-level density of states in local quantum field theory. *Communications in Mathematical Physics*, 106(2):321–344, 1986. doi: 10.1007/BF01454978. URL <https://doi.org/10.1007/BF01454978>.
- [18] J. I. Cirac and F. Verstraete. Renormalization and tensor product states in spin chains and lattices. *Journal of Physics A: Mathematical and Theoretical*, 42(50):504004, 2009. doi: 10.1088/1751-8113/42/50/504004. URL <https://doi.org/10.1088/1751-8113/42/50/504004>.
- [19] A. Cohen, I. Daubechies, and P. Vial. Wavelets on the Interval and Fast Wavelet Transforms. *Applied and Computational Harmonic Analysis*, 1:54–81, 1993. ISSN 1063-5203. doi: 10.1006/acha.1993.1005. URL <https://doi.org/10.1006/acha.1993.1005>.
- [20] R. Conti and G. Morsella. Asymptotic morphisms and superselection theory in the scaling limit II: analysis of some models. *Communications in Mathematical Physics*, 376:1767–1801, 2020. doi: 10.1007/s00220-019-03564-8. URL <https://doi.org/10.1007/s00220-019-03564-8>.
- [21] M. Cramer, A. Serafini, and J. Eisert. Locality of dynamics in general harmonic quantum systems. In M. Ericsson and S. Montangero, editors, *Quantum information and many body quantum systems*, volume 8 of *Publications of the Scuola Normale Superiore, CRM*, pages 51–72, 2008. URL <http://www.springer.com/birkhauser/mathematics/scuolanormalesuperiore/book/978-88-7642>
- [22] C. D’Antoni, G. Morsella, and R. Verch. Scaling algebras for charged fields and short-distance analysis for localizable and topological charges. *Annales Henri Poincaré*, 5:809–870, 2004. doi: 10.1007/s00023-004-0183-7. URL <https://doi.org/10.1007/s00023-004-0183-7>.
- [23] I. Daubechies. *Ten Lectures on Wavelets*, volume 61 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. SIAM, 1992. doi: 10.1137/1.9781611970104. URL <https://doi.org/10.1137/1.9781611970104>.
- [24] J. Dixmier. *C*-algebras*. North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977. ISBN 0-7204-0762-1. Translated from the French by Francis Jellet, North-Holland Mathematical Library, Vol. 15.
- [25] W. Dybalski and V. Morinelli. The Bisognano–Wichmann Property for Asymptotically Complete Massless QFT. *Communications in Mathematical Physics*, 2020. ISSN 0010-3616. doi: 10.1007/s00220-020-03755-8. URL <https://doi.org/10.1007/s00220-020-03755-8>.

- [26] D. E. Evans and Y. Kawahigashi. *Quantum symmetries on operator algebras*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1998. ISBN 0-19-851175-2. Oxford Science Publications.
- [27] G. Evenbly and G. Vidal. Algorithms for entanglement renormalization. *Physical Review B: Condensed Matter and Materials Physics*, 79(14):144108, 2009. ISSN 1098-0121. doi: 10.1103/physrevb.79.144108. URL <https://doi.org/10.1103/physrevb.79.144108>.
- [28] G. Evenbly and G. Vidal. Entanglement renormalization in free bosonic systems: real-space versus momentum-space renormalization group transforms. *New Journal of Physics*, 12(2):025007, 2010. ISSN 1367-2630. doi: 10.1088/1367-2630/12/2/025007. URL <https://doi.org/10.1088/1367-2630/12/2/025007>.
- [29] G. Evenbly and G. Vidal. Tensor Network Renormalization Yields the Multi-scale Entanglement Renormalization Ansatz. *Physical Review Letters*, 115(20):200401, Nov. 2016. ISSN 0031-9007. doi: 10.1103/physrevlett.115.200401. URL <https://doi.org/10.1103/PhysRevLett.115.200401>.
- [30] G. Evenbly and S. R. White. Entanglement Renormalization and Wavelets. *Physical Review Letters*, 116(14):140403, 2016. ISSN 0031-9007. doi: 10.1103/physrevlett.116.140403. URL <https://doi.org/10.1103/physrevlett.116.140403>.
- [31] G. Evenbly and S. R. White. Representation and design of wavelets using unitary circuits. *Physical Review A: Atomic, Molecular, and Optical Physics*, 97(5):052314, 2018. ISSN 2469-9926. doi: 10.1103/physreva.97.052314. URL <https://doi.org/10.1103/physreva.97.052314>.
- [32] R. Fernández, J. Fröhlich, and A. D. Sokal. *Random walks, critical phenomena, and triviality in quantum field theory*. Texts and Monographs in Physics. Springer, Berlin, 1992. doi: 10.1007/978-3-662-02866-7. URL <https://doi.org/10.1007/978-3-662-02866-7>.
- [33] M. E. Fisher. The renormalization group in the theory of critical behavior. *Reviews of Modern Physics*, 46(4):597–616, 1974. ISSN 0034-6861. doi: 10.1103/revmodphys.46.597. URL <https://doi.org/10.1103/RevModPhys.46.597>.
- [34] M. E. Fisher. Renormalization group theory: Its basis and formulation in statistical physics. *Reviews of Modern Physics*, 70(2):653–681, 1998. doi: 10.1103/RevModPhys.70.653. URL <https://doi.org/10.1103/RevModPhys.70.653>.
- [35] K. O. Friedrichs. *Perturbation of Spectra in Hilbert Space*, volume III of *Lectures in Applied Mathematics*. American Mathematical Society, 1965. ISBN 978-0-8218-4682-7. URL <https://bookstore.ams.org/lam-3>.
- [36] P. Fries, I. Reyes, J. Erdmenger, and H. Hinrichsen. Renormalization of lattice field theories with infinite-range wavelets. *Journal of Statistical Mechanics: Theory and Experiment*, 2019:064001, June 2018. doi: 10.1088/1742-5468/ab14d8. URL <https://doi.org/10.1088/1742-5468/ab14d8>.
- [37] J. Glimm and A. Jaffe. The $\lambda\phi_2^4$ quantum field theory without cutoffs, III. The physical vacuum. *Acta Mathematica*, 125(i):203–267, 1970. doi: 10.1007/BF02392335. URL <https://doi.org/10.1007/BF02392335>.

- [38] J. Glimm and A. Jaffe. Boson quantum field models. In *Quantum Field Theory and Statistical Mechanics*, chapter 4, pages 123–179. Birkhäuser Boston, 1985. ISBN 978-0-8176-3275-5. doi: 10.1007/978-1-4612-5158-3_6. URL https://doi.org/10.1007/978-1-4612-5158-3_6.
- [39] J. Glimm and A. Jaffe. *Quantum Field Theory and Statistical Mechanics: Expositions*. Birkhäuser Basel, 1985. ISBN 978-1-4612-5158-3. doi: 10.1007/978-1-4612-5158-3. URL <https://doi.org/10.1007/978-1-4612-5158-3>.
- [40] A. Guichardet. *Symmetric Hilbert Spaces and Related Topics*, volume 261 of *Lecture Notes in Mathematics*. Springer-Verlag Berlin Heidelberg, 1972. doi: 10.1007/bfb0070306. URL <https://doi.org/10.1007/BFb0070306>.
- [41] R. Haag. *Local Quantum Physics: Fields, Particles, Algebras*. Texts and Monographs in Physics. Springer Verlag, 2nd edition, 1996. ISBN 978-3-642-61458-3. doi: 10.1007/978-3-642-61458-3. URL <https://doi.org/10.1007/978-3-642-61458-3>.
- [42] J. Haegeman, B. Swingle, M. Walter, J. Cotler, G. Evenbly, and V. B. Scholz. Rigorous free-fermion entanglement renormalization from wavelet theory. *Physical Review X*, 8(1):011003, 2018. doi: 10.1103/PhysRevX.8.011003. URL <https://doi.org/10.1103/PhysRevX.8.011003>.
- [43] W. Härdle, G. Kerkycharian, D. Picard, and A. Tsybakov. *Wavelets, approximation, and statistical applications*, volume 129 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1997. ISBN 0-387-98453-4. doi: 10.1007/978-1-4612-2222-4. URL <https://doi.org/10.1007/978-1-4612-2222-4>.
- [44] R. Honegger and A. Rieckers. *Photons in Fock Space and Beyond*. World Scientific Publishing Company, 2015. doi: 10.1142/9251-voll. URL <https://doi.org/10.1142/9251-voll>.
- [45] A. Jaffe and J. Glimm. *Quantum Physics: A Functional Integral Point of View*. Springer, New York, 2nd edition, 1987. ISBN 978-1-4612-4728-9. doi: 10.1007/978-1-4612-4728-9. URL <https://doi.org/10.1007/978-1-4612-4728-9>.
- [46] V. F. R. Jones. A No-Go Theorem for the Continuum Limit of a Periodic Quantum Spin Chain. *Communications in Mathematical Physics*, 357(1): 295–317, Jan 2018. ISSN 1432-0916. doi: 10.1007/s00220-017-2945-3. URL <https://doi.org/10.1007/s00220-017-2945-3>.
- [47] L. P. Kadanoff. Scaling laws for Ising models near T_c . *Physics Physique Fizika*, 2(6):263, 1966. doi: 10.1103/PhysicsPhysiqueFizika.2.263. URL <https://doi.org/10.1103/PhysicsPhysiqueFizika.2.263>.
- [48] R. V. Kadison and J. R. Ringrose. *Fundamentals of the Theory of Operator Algebras, Vol. II, Advanced Theory*. Pure and Applied Mathematics. Academic Press, Inc., 1986.
- [49] T. Kato. *Perturbation Theory for Linear Operators*, volume 132 of *Die Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, New York Inc., 2nd edition, 1995. ISBN 978-3-642-66282-9. doi: 10.1007/978-3-642-66282-9. URL <https://doi.org/10.1007/978-3-642-66282-9>.

- [50] A. Kliesch and R. Koenig. Continuum limits of homogeneous binary trees and the Thompson group. *Physical Review Letters*, 124(1):010601, Jan. 2020. doi: 10.1103/PhysRevLett.124.010601. URL <https://doi.org/10.1103/PhysRevLett.124.010601>.
- [51] T. Lang, K. Liegener, and T. Thiemann. Hamiltonian Renormalisation I: Derivation from Osterwalder-Schrader Reconstruction. *Classical and Quantum Gravity*, 35(24):245011, 2018. doi: 10.1088/1361-6382/aaec56. URL <https://doi.org/10.1088/1361-6382/aaec56>.
- [52] E. H. Lieb and D. W. Robinson. The finite group velocity of quantum spin systems. *Communications in Mathematical Physics*, 28(3):425–431, Sept. 1972. doi: 10.1007/BF01645779. URL <https://doi.org/10.1007/BF01645779>.
- [53] R. Longo. A simple proof of the existence of modular automorphisms in approximately finite-dimensional von Neumann algebras. *Pacific J. Math.*, 75(1):199–205, 1978. ISSN 0030-8730. URL <http://projecteuclid.org/euclid.pjm/1102810157>.
- [54] R. Longo and F. Xu. Von Neumann Entropy in QFT. *Preprint, arXiv: 1911.09390*, 2019. URL <http://arxiv.org/abs/1911.09390>.
- [55] R. Longo, V. Morinelli, F. Preta, and K.-H. Rehren. Split property for free massless finite helicity fields. *Ann. Henri Poincaré*, 20(8):2555–2584, 2019. ISSN 1424-0637. doi: 10.1007/s00023-019-00820-4. URL <https://doi.org/10.1007/s00023-019-00820-4>.
- [56] Masamichi Takesaki. *Theory of operator algebras. III*, volume 127 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. ISBN 3-540-42913-1. doi: 10.1007/978-3-662-10453-8. URL <https://doi.org/10.1007/978-3-662-10453-8>. Operator Algebras and Non-commutative Geometry, 8.
- [57] H. Matsueda. Analytic Optimization of a MERA network and its Relevance to Quantum Integrability and Wavelet. *Preprint, arXiv: 1608.02205*, 2016. URL <https://arxiv.org/abs/1608.02205>.
- [58] Y. Meyer. *Wavelets and operators*, volume 37 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 1989. doi: 10.1017/CBO9780511623820. URL <https://doi.org/10.1017/CBO9780511623820>.
- [59] A. Milsted and T. J. Osborne. Quantum Yang-Mills theory: An overview of a program. *Physical Review D: Particles and Fields*, 98(1):014505, 2018. ISSN 2470-0010. doi: 10.1103/physrevd.98.014505. URL <https://doi.org/10.1103/PhysRevD.98.014505>.
- [60] V. Morinelli. The Bisognano-Wichmann property on nets of standard subspaces, some sufficient conditions. *Ann. Henri Poincaré*, 19(3):937–958, 2018. ISSN 1424-0637. doi: 10.1007/s00023-017-0636-4. URL <https://doi.org/10.1007/s00023-017-0636-4>.
- [61] B. Nachtergaele and R. Sims. Lieb-Robinson bounds in quantum many-body physics. In D. U. Robert Sims, editor, *Entropy and the Quantum*, volume 529 of *Contemporary Mathematics*, pages 141–176. American Mathematical Society, 2010. ISBN 978-0-8218-5247-7. URL <https://arxiv.org/abs/1004.2086>.

- [62] B. Nachtergaele, H. Raz, B. Schlein, and R. Sims. Lieb-Robinson bounds for harmonic and anharmonic lattice systems. *Communications in Mathematical Physics*, 286(3):1073–1098, 2009. doi: 10.1007/s00220-008-0630-2. URL <https://doi.org/10.1007/s00220-008-0630-2>.
- [63] B. Nachtergaele, B. Schlein, R. Sims, S. Starr, and V. Zagrebnov. On the existence of the dynamics for anharmonic quantum oscillator systems. *Reviews in Mathematical Physics*, 22(02):207–231, 2010. doi: 10.1142/S0129055X1000393X. URL <https://doi.org/10.1142/S0129055X1000393X>.
- [64] B. Nachtergaele, R. Sims, and A. Young. Quasi-locality bounds for quantum lattice systems. I. Lieb-Robinson bounds, quasi-local maps, and spectral flow automorphisms. *Journal of Mathematical Physics*, 60(6):061101, 2019. ISSN 0022-2488. doi: 10.1063/1.5095769. URL <https://doi.org/10.1063/1.5095769>.
- [65] T. J. Osborne. Continuum Limits of Quantum Lattice Systems. *Preprint, arXiv:1901.06124*, 2019. URL <http://arxiv.org/abs/1901.06124v1>.
- [66] T. J. Osborne and D. E. Stiegemann. Quantum fields for unitary representations of Thompson’s groups F and T. *Preprint, arXiv:1903.00318*, 2019. URL <http://arxiv.org/abs/1903.00318v1>.
- [67] R. N. C. Pfeifer, G. Evenbly, and G. Vidal. Entanglement renormalization, scale invariance, and quantum criticality. *Physical Review A: Atomic, Molecular, and Optical Physics*, 79(4):040301, 2009. ISSN 1050-2947. doi: 10.1103/physreva.79.040301. URL <https://doi.org/10.1103/PhysRevA.79.040301>.
- [68] M. Reed and B. Simon. *Methods of Modern Mathematical Physics: Vol.: 2.: Fourier Analysis, Self-Adjointness*. Academic Press, 1975. URL <https://www.elsevier.com/books/ii-fourier-analysis-self-adjointness/reed/978-0-08-092>.
- [69] S. Sachdev. *Quantum Phase Transitions*. Cambridge University Press, second edition, 2011. ISBN 9780511973765. doi: 10.1017/CBO9780511973765. URL <https://doi.org/>.
- [70] U. Schollwöck. The density-matrix renormalization group. *Reviews of Modern Physics*, 77:259–315, 2005. ISSN 0034-6861. doi: 10.1103/revmodphys.77.259. URL <https://doi.org/10.1103/revmodphys.77.259>.
- [71] I. E. Segal. Mathematical characterization of the physical vacuum for a linear Bose-Einstein field. *Illinois Journal of Mathematics*, 6(3):500–523, 1962. doi: 10.1215/ijm/1255632508. URL <https://projecteuclid.org/euclid.ijm/1255632508>.
- [72] E. M. Stein and G. Weiss. *Introduction to Fourier analysis on Euclidean spaces*. Number 32 in Princeton Mathematical Series. Princeton University Press, Princeton, N.J., 1972. doi: 10.1515/9781400883899. URL <https://doi.org/10.1515/9781400883899>.
- [73] A. Stottmeister, V. Morinelli, G. Morsella, and Y. Tanimoto. Operator-algebraic renormalization and wavelets. *Preprint, arXiv: 2002.01442*, 2020. URL <https://arxiv.org/pdf/2002.01442v1>.
- [74] T. Thiemann. Canonical Quantum Gravity, Constructive QFT and Renormalisation. *Preprint, arXiv: 2003.13622*, 2020. URL <https://arxiv.org/pdf/2003.13622v1>.

- [75] G. Vidal. A Class of Quantum Many-Body States That Can Be Efficiently Simulated. *Physical Review Letters*, 101(11):110501, Sept. 2008. ISSN 0031-9007. doi: 10.1103/physrevlett.101.110501. URL <https://doi.org/10.1103/PhysRevLett.101.110501>.
- [76] F. J. Wegner. Corrections to scaling laws. *Physical Review B*, 5(11):4529, 1972. doi: 10.1103/PhysRevB.5.4529. URL <https://doi.org/10.1103/PhysRevB.5.4529>.
- [77] S. R. White. Density matrix formulation for quantum renormalization groups. *Physical review letters*, 69(19):2863, 1992. doi: 10.1103/PhysRevLett.69.2863. URL <https://doi.org/10.1103/PhysRevLett.69.2863>.
- [78] S. R. White. Density-matrix algorithms for quantum renormalization groups. *Physical Review B*, 48(14):10345, 1993. doi: 10.1103/PhysRevB.48.10345. URL <https://doi.org/10.1103/PhysRevB.48.10345>.
- [79] S. R. White and R. M. Noack. Real-space quantum renormalization groups. *Physical Review Letters*, 68(24):3487, 1992. doi: 10.1103/PhysRevLett.68.3487. URL <https://doi.org/10.1103/PhysRevLett.68.3487>.
- [80] K. G. Wilson. Renormalization Group and Critical Phenomena. I. Renormalization Group and the Kadanoff Scaling Picture. *Physical Review B: Condensed Matter and Materials Physics*, 4(9):3174–3183, 1971. ISSN 0556-2805. doi: 10.1103/physrevb.4.3174. URL <https://doi.org/10.1103/physrevb.4.3174>.
- [81] K. G. Wilson. Renormalization Group and Critical Phenomena. II. Phase-Space Cell Analysis of Critical Behavior. *Physical Review B: Condensed Matter and Materials Physics*, 4(9):3184–3205, 1971. ISSN 0556-2805. doi: 10.1103/physrevb.4.3184. URL <https://doi.org/10.1103/physrevb.4.3184>.
- [82] K. G. Wilson. The renormalization group: Critical phenomena and the Kondo problem. *Reviews of Modern Physics*, 47(4):773, 1975. doi: 10.1103/RevModPhys.47.773. URL <https://doi.org/10.1103/RevModPhys.47.773>.
- [83] F. Witteveen and M. Walter. Wavelet construction of bosonic entanglement renormalization circuits. *Preprint, arXiv: 2004.11952*, 2020. URL <https://arxiv.org/pdf/2004.11952v1>.
- [84] F. Witteveen, V. Scholz, B. Swingle, and M. Walter. Quantum circuit approximations and entanglement renormalization for the Dirac field in 1+1 dimensions. *Preprint, arXiv: 1905.08821*, 2019. URL <https://arxiv.org/pdf/1905.08821v1>.
- [85] J. Yngvason. A note on essential duality. *Lett. Math. Phys.*, 31(2):127–141, 1994. ISSN 0377-9017. doi: 10.1007/BF00750147. URL <https://doi.org/10.1007/BF00750147>.
- [86] M. S. Zini and Z. Wang. Conformal Field Theories as Scaling Limit of Anyonic Chains. *Communications in Mathematical Physics*, 363(3):877–953, 2018. doi: 10.1007/s00220-018-3254-1. URL <https://doi.org/10.1007/s00220-018-3254-1>.
- [87] J. Zinn-Justin. *Phase Transitions and Renormalization Group*. Oxford University Press, 2007. ISBN 9780199227198. doi: 10.1093/acprof:oso/9780199227198.001.0001. URL <https://doi.org/10.1093/acprof:oso/9780199227198.001.0001>.